

**APPLICATION FOR  
UNITED STATES PATENT**

under

37 CFR §1.53(b)

**For**

**APOPTOSIS MODULATOR BCL-B AND METHODS FOR MAKING AND USING  
SAME**

Applicants:

John C. Reed

Ning Ke

Adam Godzik

Assignee: The Burnham Institute, San Diego, CA

ATTORNEY DOCKET:

DATE OF DEPOSIT:

February 7, 2002

EXPRESS MAIL NO.:

EL 754073637 US

**APOPTOSIS MODULATOR BCL-B AND METHODS FOR MAKING**  
**AND USING SAME**

**PRIORITY INFORMATION**

This application claims priority to U.S. application serial no. 60/267,166, filed

5 February 7, 2001.

**TECHNICAL FIELD**

This invention generally relates to cell and molecular biology and the regulation of cell proliferation, apoptosis and survival. In particular, the invention provides polypeptides comprising apoptosis modulator Bcl-B, a Bcl-2 family member, nucleic acids encoding the polypeptides, and methods for making and using these compositions, including, for example, modulating cell apoptosis, survival, proliferation.

**BACKGROUND**

Programmed cell death or apoptosis is a cellular suicide process in which damaged or harmful cells are eliminated from multicellular organisms. Cells undergoing apoptosis have distinct morphological changes including cell shrinkage, membrane blebbing, chromatin condensation, apoptotic body formation and protein and nucleic acid fragmentation. This cellular suicide program is evolutionarily conserved across animal and plant species.

Apoptosis plays an important role in the development and homeostasis of metazoans and is also important for insect embryonic development and metamorphosis. Furthermore, apoptosis can act as a host defense mechanism. For example, apoptosis eliminates virally infected cells thereby limiting propagation of viruses. Apoptosis is also involved in plant reactions to biotic and abiotic insults. Moreover, dysregulation of apoptosis has been associated with a variety of human diseases including cell proliferative disorders (*e.g.*, cancer), cell degenerative disorders (*e.g.*, neurodegeneration, muscular degeneration, ischemia, stroke, etc.) and autoimmune diseases. Accordingly, identification of the components that modulate apoptosis provides a means to study and manipulate the process in a wide variety of organisms.

Programmed cell death is regulated by the interplay of proteins that inhibit and proteins that stimulate cell death or cell survival. Among the proteins that modulate apoptosis are the Bcl-2 family members. Bcl-2 protein family members include proteins that promote and inhibit

programmed cell death. Bcl-2 family proteins play a role in apoptosis regulation in metazoan species. In humans, over 20 Bcl-2 proteins have been identified to date, including proteins which suppress (Bcl-2, Bcl-XL, Mcl-1, Bfl-1/A1, Bcl-W) and proteins which promote (Bax, Bak, Bok, Bad, Bid, Bik, Bim, Nip3, Nix) cell death (Reed, J. *Oncogene* 17, 3225-3236(1998); Reed, J. C. *Amer J Pathol* 157, 1415-1430(2000)).

Bcl-2 family proteins contain at least one of four conserved regions, termed Bcl-2 Homology (BH) domains. Most members of this family also contain a transmembrane (TM) domain located near the carboxyl-terminus that anchors them in intracellular membranes of mitochondria and other organelles (Reed, J. *Oncogene* 17, 3225-3236(1998); Reed, J. C. *Amer J Pathol* 157, 1415-1430(2000)).

Many Bcl-2 family proteins are capable of physically interacting, forming homo- or hetero-dimers, and functioning as agonists or antagonists of each other (Reed, J. *Oncogene* 17, 3225-3236 (1998); Reed, J. C. *Amer J Pathol* 157, 1415-1430 (2000); Oltvai, Z. N., and Korsmeyer, S. J. *Cell* 79, 189-192 (1994)). Specificity for interaction partners and tissue-specific patterns of expression combine to endow each Bcl-2 protein with a physiological role *in vivo*, resulting for example in highly diverse phenotypes when members of this multigene family are individually knocked-out in mice (Vaux, D. and Korsmeyer, S. *Cell* 96, 245-254 (1999)).

Thus, a need exists to identify members of the Bcl-2 family and to elucidate their functional characteristics. The present invention we describe the molecular cloning and initial characterization of a new human member of the Bcl-2 family, Bcl-B.

## SUMMARY

The present invention is based in part on the identification and characterization of a novel member of the Bcl-2 family of apoptosis modulators, denoted Bcl-B. Bcl-B is capable of modulating apoptosis in cells. For example, Bcl-B inhibits apoptosis induced by Bax. Bcl-B also binds to itself as well as other modulators of apoptosis including, for example, Bax, Bcl-2 and Bcl-XL. Thus, Bcl-B is involved in apoptotic signaling as well as modulating activity or activation of other proteins, or having its own activity modulated by other proteins associated with programmed cell death. Accordingly, compositions of the invention, including, for example, Bcl-B polypeptides, polynucleotides, antibodies and subsequences thereof are useful for modulating apoptosis and associated signaling pathways, as well as for detecting Bcl-B (*e.g.*,

for diagnosis or monitoring Bcl-B based therapy) and identifying agents that affect Bcl-B expression, activity or binding.

Thus, in accordance with the invention there are provided isolated and recombinant Bcl-B nucleic acids. In one embodiment, a Bcl-B nucleic acid includes a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1, wherein the sequence is distinct from EST Accession no. AA098865. In additional embodiments, a Bcl-B nucleic acid includes a polynucleotide sequence having at least about 80%, 90%, 95% or more identity to SEQ ID NO:1, wherein the sequence is distinct from EST Accession no. AA098865. In still other embodiments, a Bcl-B nucleic acid encodes a polypeptide that modulates apoptosis (e.g., a sequence set forth in or including SEQ ID NO:2). Polynucleotide sequences included in the invention can be of any size, for example, a sequence less than about 50 kB, less than about 25 kB, less than about 10 kB, less than about 5 kB, less than about 2.5 kB, or between about 2.5 kB and 1 kB, 1 kB and 0.5 kB, 0.5 kB and 0.25 kB and 0.1kB and 15 base pairs.

In additional embodiments, isolated or recombinant nucleic acids include SEQ ID NO:1; SEQ ID NO:1, wherein one or more T's are U; nucleic acid sequences complementary to SEQ ID NO:1 and SEQ ID NO:1, wherein one or more T's are U; and subsequences thereof that are at least 15 base pairs long, e.g., a length of about 12-30, 30-50, 50-100, 100-250, 250-500, 500-1000, 1000-2500, 2500-5000 or 5000-10000 base pairs.

In yet additional embodiments, nucleic acids include sequences that hybridize to a sequence set forth as SEQ ID NO:1, wherein the nucleic acid is distinct from Accession no. AA098865. In various aspects, the sequences hybridize under low, moderate or stringent hybridization conditions.

Invention nucleic acids can be expressed (e.g., transcribed and/or translated) in solution or solid phase, or in cells tissue or organs *in vitro*, *ex vivo* or *in vivo*. In order to express the Bcl-B polynucleotides, they can be linked to cis-acting nucleic acid sequences, such as promoters, enhancers and other transcription/translation control elements. Thus, the invention further provides expression cassettes including a Bcl-B polynucleotide sequence operably linked to an expression control element. In one embodiment, a polynucleotide sequence operably linked to an expression control element has at least about 70% identity to SEQ ID NO:1. In one aspect, the polynucleotide sequence encodes a polypeptide that inhibits apoptosis or an antisense that stimulates or induces apoptosis (e.g., stimulates or inhibits Bax mediated apoptosis). In another

aspect, the polynucleotide sequence encodes a polypeptide comprising SEQ ID NO:2. In other embodiments, the expression cassette includes a promoter or enhancer, for example, a constitutive, inducible, tissue-specific or developmentally regulated promoter or enhancer. In additional embodiments, an expression cassette further comprises a vector. In various aspects, a vector confers expression in bacteria, plant, insect, mammalian or a yeast cell. In additional aspects, a vector comprises a viral vector such as an adenovirus, retrovirus, adenovirus, adeno-associated virus, lentivirus, reovirus, rotavirus, herpes simplex virus, parvovirus, papilloma virus or cytomegalovirus.

The invention also provides isolated and recombinant Bcl-B polypeptides. In one embodiment, an isolated or recombinant polypeptide includes a sequence having at least about 65% identity to SEQ ID NO:2, and having one or more activities of the polypeptide set forth in SEQ ID NO:2. In various aspects, the isolated or recombinant polypeptide has at least about 75%, 85%, 90%, 95% or more identity to SEQ ID NO:2. Polypeptide sequences included in the invention can be of any size, for example, a sequence at least about 15, 20, 25, 30, 40, 50, 75, 125, 150 or 200 or more amino acids in length. Bcl-B polypeptides generally have one or more activities as set forth herein, for example, modulating apoptosis (e.g., inhibits Bax mediated apoptosis) in a cell (e.g., bacteria, plant, insect, mammalian or yeast). In various aspects, the cell is a human cell such as a heart, brain, lung, kidney, liver, pancreas, spleen, thymus, colon, leukocyte, small intestine, testis, prostate or ovarian cell, and optionally expresses bax.

Polypeptide sequences of the invention include modified forms that retain, at least a part of, one or more activities or functions of wild type Bcl-B. Exemplary Bcl-B activities include, for example, modulating apoptosis, homodimerization, heteromerization, binding to Bcl-2, Bcl-XL or Bax, forming a membrane channel, associating with mitochondria, or immunogenicity. In one embodiment, the polypeptide contains a transmembrane domain substantially the same as the underlined amino acid sequence set forth in Figure 1A (SEQ ID NO:13). In another embodiment, the polypeptide contains one or more BH1, BH2, BH3 or BH4 domains set forth in Figure 1A.

Polypeptide sequences of the invention include chimeric forms of Bcl-B (e.g., having at least about 65% identity to SEQ ID NO:2, optionally encoded by a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1) and fused to a second polypeptide sequence (e.g., different amino acid sequences). In one embodiment, a second polypeptide sequence

comprises a tag. In another embodiment, a Bcl-B polypeptide includes a transmembrane domain of a mitochondrial protein or another Bcl-2 protein family member transmembrane domain.

The invention further provides antibodies that specifically bind to a Bcl-B polypeptide. In one embodiment, an antibody binds to a sequence in or including a sequence set forth in SEQ ID NO:2, for example, an immunogenic subsequence thereof. In other embodiments, the antibody modulates an activity or function of Bcl-B. Exemplary Bcl-B activities modulated by an antibody include, for example, one or more of modulating apoptosis, homodimerization, heteromerization, binding to Bcl-2, Bcl-XL or Bax, forming a membrane channel, associating with mitochondria, or immunogenicity.

Invention nucleic acids, polypeptides, antibodies and other compositions set forth herein may be attached to a substrate. Thus, the invention provides substrates that include Bcl-B nucleic acids, polypeptides and antibodies suitable for detection, genetic or expression profiling or diagnosis. For example, in one embodiment, a Bcl-B nucleic acid, polypeptide or antibody is attached to a two-dimensional detection substrate in which a plurality of other molecules are attached at discrete defined positions (i.e., addresses).

Invention nucleic acids, polypeptides, antibodies and other compositions set forth herein may be expressed in cells, *in vitro*, *ex vivo* or *in vivo*. Thus, the invention provides transformed cells including a Bcl-B nucleic acid, polypeptide or antibody. In various aspects, the cell is a bacteria, plant, insect, mammalian (e.g., human) or yeast cell.

Invention nucleic acids, polypeptides, antibodies and other compositions set forth herein may be expressed in animals, including non-human transgenic animals. In one embodiment, a non-human transgenic animal includes a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1. In another embodiment, a non-human transgenic animal expresses a polypeptide or an antisense that modulates apoptosis. In yet another embodiment, expression of the polypeptide or antisense is tissue-specific or is in one or more cells of a non-human transgenic animal (e.g., in one or more of heart, brain, lung, kidney, liver, pancreas, spleen, thymus, colon, muscle, leukocyte, small intestine, testis, prostate or ovary). Transgenic animals include progeny thereof, homozygous or heterozygous in respect to the Bcl-B molecule.

Invention nucleic acids, polypeptides, antibodies and other compositions set forth herein may be expressed in plants. The invention therefore provides plants, plant parts and seeds that can express Bcl-B nucleic acids, polypeptides and antibodies. In one embodiment, a transgenic

plant, plant part or seed includes a nucleic acid sequence having at least about 70% identity to SEQ ID NO:1. In another embodiment, a transgenic plant, plant part or seed includes a nucleic acid encoding a polypeptide that modulates apoptosis. In one aspect, at least a portion of the plant exhibits a decreased level of senescence. In yet another embodiment, a transgenic plant,  
5 plant part or seed is resistant to abiotic insult, such as an insult induced by high moisture, low moisture, salinity, nutrient deficiency, air pollution, high temperature, low temperature, soil toxicity, herbicide or insecticide, or biotic insult, such as an insult induced by a plant pathogen (e.g., a virus, fungus, bacteria or nematode).

Invention nucleic acids, polypeptides, antibodies, including cells, plants, substrates, transgenic animals and other compositions set forth herein may be included in kits. Thus, the invention provides kits including, *inter alia*, Bcl-B nucleic acids, polypeptides, antibodies, including cells, plants, substrates and transgenic animals in a container. In various embodiments, a kit includes instructions suitable for using one or more of the kit components. In one aspect, instructions are for detecting a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1, a polynucleotide or polypeptide that binds to a polypeptide of claim 46, or a polypeptide in or including a sequence set forth in SEQ ID NO:2.

Invention nucleic acids, polypeptides, antibodies, including cells, may be included in pharmaceutical formulations. Thus, the invention additionally provides isolated or recombinant Bcl-B nucleic acids, polypeptides, antibodies and cells in a pharmaceutically acceptable carrier.  
20 Pharmaceutical carriers include those suitable for particular routes of administration including, for example, intracranial, intravenous, intramuscular, subcutaneous, via intubation, inhalation, oral, topical (ocular or nasal), or intra-cavity (e.g., rectal or vaginal).

The invention furthermore provides methods of producing Bcl-B nucleic acids, polypeptides, antibodies, including substrates, cells, plants, transgenic animals and other  
25 compositions set forth herein. In one embodiment, a method of producing a Bcl-B polypeptide includes expressing a nucleic acid encoding an amino acid sequence having at least about 65% identity to SEQ ID NO:2 in solution, in solid phase, or in a cell *in vitro* or *in vivo*.

The invention moreover provides methods for detecting Bcl-B nucleic acids and polypeptides, including cells, plants, transgenic animals and other compositions containing Bcl-B  
30 B set forth herein. In one embodiment, a method for detecting the presence of a polynucleotide sequence encoding a Bcl-B amino acid sequence having at least about 65% identity to SEQ ID

NO:2, or an encoded Bcl-B polypeptide, includes: contacting a sample with a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1 or a Bcl-B antibody; and detecting the presence of a polynucleotide sequence encoding an amino acid sequence having at least about 65% identity to SEQ ID NO:2, or a Bcl-B polypeptide in the sample.

5 Invention compositions including Bcl-B nucleic acids (sense and antisense), polypeptides and antibodies are useful for modulating apoptosis of cells *in vitro*, *in vivo* and *ex vivo*. Thus, the invention provides methods for modulating apoptosis of cells, *in vitro*, *in vivo* and *ex vivo*, optionally in a subject, such as a mammalian (e.g., human) subject. In one embodiment, a method for modulating apoptosis includes contacting a cell with a Bcl-B polypeptide, a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1 or an antisense thereof, or a Bcl-B antibody in an amount sufficient to modulate apoptosis in the cell. In one aspect, the antisense comprises a sequence complementary to Bcl-B sense strand, a sequence that forms a triplex with Bcl-B, a ribozyme, a DNAzyme or an RNAi molecule. In another aspect, apoptosis is induced or increased. In another aspect, apoptosis is prevented or inhibited. In yet another aspect, the cell is at risk of apoptosis or is undergoing apoptosis. In a further aspect, the cell is or is at risk of undesirable proliferation or is undergoing hyperproliferation. In still another aspect, the cell expresses Bax. Exemplary cells in which apoptosis can be modulated include, for example, heart, brain, lung, kidney, liver, pancreas, spleen, thymus, colon, leukocyte, small intestine, testis, prostate and ovarian cells.

20 Methods of the invention further include modulating apoptosis in a subject. In one embodiment, a method includes administering to the subject an amount of a Bcl-B polypeptide, a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1 or an antisense thereof, or a Bcl-B antibody sufficient to modulate apoptosis in the subject. Candidate subjects having or at risk of a disorder or pathological condition associated with apoptosis may be treated. Thus in another embodiment, a method includes administering to a subject having or at risk of a disorder associated with apoptosis with an amount of a Bcl-B polypeptide, a polynucleotide sequence having at least about 70% identity to SEQ ID NO:1 or an antisense thereof, or a Bcl-B antibody sufficient to treat the subject having or at risk of a disorder associated with apoptosis. In one aspect, the antisense comprises a sequence complementary to Bcl-B sense strand, a sequence that forms a triplex with Bcl-B, a ribozyme, a DNAzyme or an RNAi molecule.

30



Subjects suffering or at risk of suffering from a disorder associated with apoptosis include those having a cell degenerative or proliferative disorders. Thus, disorders treatable in a method of the invention include, for example, a neoplasia, autoimmune disorder or fibrotic condition. In particular aspects, the disorder is characterized by neural or muscle degeneration, such as Alzheimer's disease, Parkinson's disease, Creutzfeldt-Jacob's disease (CJD), Huntington disease (HD), Machado-Joseph disease (MJD), Spinocerebellar ataxias 1, 2 and 6 (SCA-1, -2 and -6), dentatorubropallidolusian atrophy (DRPLA), Kennedy's disease, ischemia, stroke and head trauma, for example. In additional particular aspects, apoptosis or the disorder associated with apoptosis is present in one or more cells of heart, brain, lung, kidney, liver, pancreas, spleen, thymus, colon, muscle, leukocyte, small intestine, testis, prostate or ovary.

Invention compositions are useful as reagents. For example, as Bcl-B modulates apoptosis, genes or agents or molecules that modulate Bcl-B expression, activity or function, for example, promote or inhibit Bcl-B expression or Bcl-B binding with bax, in turn modulate apoptosis. Thus, the invention also provides methods for identifying a gene or agent or molecule in solution, in solid phase, or in a cell that modulates expression or activity of a Bcl-B polypeptide. In one embodiment, a method includes: contacting a cell that expresses Bcl-B polypeptide with a test gene or test agent; and measuring expression or activity of Bcl-B or nucleic acid encoding Bcl-B polypeptide, wherein an increase or decrease in the amount of Bcl-B polypeptide or nucleic acid encoding Bcl-B polypeptide or activity of Bcl-B polypeptide identifies the test gene or test agent as a modulator of Bcl-B polypeptide's expression or activity. In another embodiment, a method includes contacting Bcl-B with a test agent and identifying an agent (e.g., polypeptide) that binds to Bcl-B. In one aspect, Bcl-B polypeptide contains one or more 15N-labeled amino acids, and binding is detected by resonance changes in Bcl-B polypeptide. In another aspect, the test agent (i.e., molecule) comprises a polypeptide sequence (e.g., an antibody). In yet another aspect, the test agent is attached to the surface of a substrate. As set forth herein, test agents include libraries of molecules. Test agents may be attached at discrete positions of a substrate. The binding agent identified can subsequently be tested for modulating one or more activities or functions of Bcl-B.

In yet another embodiment, a method includes contacting Bcl-B with a binding molecule (e.g., Bax, Bcl-B, or a Bcl-2 protein family member such as Bcl-2 or Bcl-XL) under conditions allowing binding in the presence and absence of a test agent; and measuring binding between

1  
2  
3  
4  
5 Bcl-B and the molecule (e.g., Bax, Bcl-B, or a Bcl-2 protein family member such as Bcl-2 or Bcl-XL) in the presence and absence of the test agent; increased or decreased binding identifies an agent that modulates binding of Bcl-B to the binding molecule. In various aspects, the method is performed in solution, solid phase or in a cell, *in vitro* or *in vivo*, for example, in a bacteria, plant, insect, mammalian or yeast cell. In another aspect, the assay comprises a two-hybrid system for expressing Bcl-B and the binding molecule. In another aspect, *in vitro* binding is measured by detecting fluorescence of Bcl-B conjugated to a fluorophore.

6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211

sample. Samples may be from a subject, in which case they may detect abnormal amounts of Bcl-B which in turn indicates a disorder associated with apoptosis. Thus, in various aspects, the sample is obtained from a subject having or at risk of having a cell proliferative (e.g., hyperproliferation) or degenerative disorder (e.g., undesirable apoptosis).

## DESCRIPTION OF DRAWINGS

**Figure 1** shows (A) The predicted amino-acid sequence of exemplary Bcl-B (SEQ ID NO:2); the transmembrane domain (TM) is underlined. (B) An alignment of Bcl-B BH1, BH2, BH3 and BH4 domains with other Bcl-2 family proteins (SEQ ID NOs:13-36). Numbers on the left indicate the position of the amino-acid in each protein based on Genbank AAD08703 (murine Bcl-2), Q90343 (chicken Bcl-2), AAA35591 (hu Bcl-2), CAA80661 (hu Bcl-XL), and P41958 (*C.elegans CED9*); black and gray boxes indicate identical and similar residues, respectively; asterisks under the BH2 alignment indicate the intron junction for hu *BCL-B*, *BCL-2*, and *BCL-XL* genes. (C) Expression of BCL-B in adult human tissues.

**Figures 2A-2E** show that Bcl-B interacts with itself and several Bcl-2 family proteins. Expressed chimeras Myc-Bcl-B, Myc-hCBP, GFP, GFP-Bcl-B, GFP-Bak, Bcl-2, Flag-Bcl-XL, HA-Bax, HA-BAG1 and Flag-Bcl-B were immunoprecipitated (IP) with anti-Myc or anti-Flag monoclonal antibodies and western blotted (WB) with antibodies specific for GFP, HA, Flag, Myc, or Bcl-XL.

**Figure 3** shows that Bcl-B inhibits Bax- but not Bak-induced apoptosis in (A) 293T and (B) Cos-7 cells. The percentage of GFP-positive cells with apoptotic morphology (fragmented nuclei or condensed chromatin) was determined (mean  $\pm$  SD; n = 3).

**Figure 4** shows that Bcl-B binding with Bax correlates with protection from apoptosis. (A) 293T cells transiently transfected with Myc-tagged constructs (human calcyclin-binding protein, wildtype Bcl-B, L86A, R96Q, FFR(169-171)AAA,  $\Delta$ 118-133), immunoprecipitated with anti-Myc antibody and subject to immunoblot analysis using antibodies specific for Bax. (B) caspase assays of 293T cells co-transfected with Bax and the Bcl-B constructs. Enzyme activity was determined by the release of AFC-fluorescence.

**Figure 5** shows that the C-terminal transmembrane (TM) domain of Bcl-B mediates efficient membrane targeting and is important for modulating apoptosis. (A) Confocal microscopy analysis of GFP-Bcl-B transfected Cos-7 cells. (B, C, D) 293T cells were

transfected with (B) Myc-Bcl-B, (C) Bcl-2, and (D) Myc-Bcl-B or Myc-Bcl-BATM. (E) 293T cells co-transfected with GFP (used as a marker for transfection) and either pcDNA3 (control; CNTL) or pcDNA3-Bax, in combination with a 2-fold excess of pcDNA3 (control), pcDNA3Myc-Bcl-B or pcDNA3-Myc-Bcl-BATM. The percentage of green cells with apoptotic morphology was determined (mean  $\pm$  SD; n = 3).

## DETAILED DESCRIPTION

The invention provides polypeptides that “modulate apoptosis,” proteins that increase or induce apoptosis or decrease or prevent programmed cell death, including exemplary Bcl-B (Figure 1A, *e.g.*, SEQ ID NO:2). The invention also provides nucleic acids encoding polypeptides that “modulate apoptosis,” including exemplary Bcl-B (Example 2, *e.g.*, SEQ ID NO:1). The invention further provides antibodies that bind to Bcl-B including antibodies that increase an activity of Bcl-B (*i.e.*, an agonistic antibody) and antibodies that decrease an activity of Bcl-B (*i.e.*, an antagonistic antibody). Also provided are polynucleotides (*e.g.*, oligonucleotide probes) of various lengths that hybridize to Bcl-B set forth in SEQ ID NO:1. Methods for making these compositions are also provided. Kits containing invention compositions in a container, including instructions for use (*e.g.*, written on a hard copy or a computer readable medium), for example, for modulating apoptosis, for detecting Bcl-B activity, expression or binding, or identifying an agent that modulates Bcl-B activity, expression or binding are provided.

Invention compositions have various activities including the ability to modulate apoptosis. Thus, invention compositions are useful as therapeutic agents for increasing or decreasing apoptosis. For example, Bcl-B protein, having wild type function or dominant negative activity, can be expressed in a cell, tissue or organ in order to modulate apoptosis. Antisense and antisense like molecules such as ribozymes, DNazymes or RNAi (RNA interference) molecules (collectively referred to herein as “antisense”) that inhibit expression of Bcl-B can reduce the level of Bcl-B gene expression, thus reducing the level of Bcl-B activity. Accordingly, the compositions can be used to manipulate apoptotic (“cell death”) mechanisms in a variety of cell types, including insect, plant and mammalian, such as human, cells, and organisms. The invention therefore provides methods for increasing or decreasing apoptosis in a

cell, tissue or organ *in vitro*, *ex vivo* or *in vivo* (e.g., in a subject in need of increased or decreased apoptosis).

Invention compositions also are useful as diagnostic markers. In particular, Bcl-B modulates apoptosis and, therefore, may indicate normal or altered apoptosis. For example, greater than normal levels of Bcl-B in a cell, tissue or organ may characterize the cell, tissue or organ as proliferative or hyperproliferative. In contrast, less than normal levels of Bcl-B in a cell, tissue or organ B may characterize the cell, tissue or organ as slowly proliferating, non-proliferating, at risk of apoptosis or undergoing apoptosis. Thus, detecting Bcl-B (nucleic acid or protein) or Bcl-B activity or expression can be used to as a diagnostic marker. Detecting Bcl-B activity or expression may also be useful for assessing a subject's risk of developing altered apoptosis or indicate a prognosis for a condition or disorder associated with apoptosis.

Bcl-B compositions are also useful for identifying agents, including therapeutic agents for treating disorders associated with apoptosis. In a method of the invention, Bcl-B (protein or nucleic acid) is contacted with a test agent and binding of Bcl-B to the agent is determined. Binding of Bcl-B to binding molecules (e.g., Bax, Bcl-2 or Bcl-X<sub>L</sub>) in the presence of a test agent can be used to identify agents that inhibit or promote Bcl-B binding to a binding molecule. In another method of the invention, Bcl-B activity or expression is determined in the presence of a test agent in order to identify agents that modulate Bcl-B activity or expression. Agents that modulate Bcl-B activity, expression or binding to a binding molecule are useful for modulating apoptosis. Agents that bind Bcl-B may also be useful as markers to indicate the presence of Bcl-B.

Additional invention methods are based on other Bcl-B activities, such as the ability of Bcl-B to bind to cellular molecules associated with apoptosis. Invention compositions are therefore also useful as therapeutic targets. For example, in a method of the invention, contacting Bcl-B with a molecule that binds Bcl-B, for example, contacting Bcl-B with Bax, Bcl-2 or Bcl-X<sub>L</sub> can be used to modulate a Bcl-B activity, such as apoptosis (increasing or decreasing). Bcl-B can also be targeted with agents that modulate Bcl-B activity or expression (e.g., antisense or ribozymes or DNazymes or RNAi), or Bcl-B binding to binding molecules.

Modulating apoptosis in cells with invention compositions allows treatment of physiological conditions associated with undesirable or insufficient apoptosis, or undesirable cell survival, growth or proliferation (e.g., where apoptosis is less than normal or desired), *ex vivo* or

*in vivo* in a subject. Conditions associated with undesirable apoptosis/cell death or undesirable cell survival/growth/proliferation include cell degenerative and proliferative disorders. Organ and organ systems affected by degenerative disorders include, for example, the nervous system (brain, spinal cord, neurons associated with tissues and organs) skeletal-muscular system (voluntary muscles) and the cardio-respiratory system (heart and blood vessels). Specific examples of neurodegeneration include cell death caused by stroke and head trauma, Alzheimer's disease, Parkinson's disease, Creutzfeldt-Jacob's disease (CJD), Huntington disease (HD), Machado-Joseph disease (MJD), ataxias (Spinocerebellar ataxias 1, 2 and 6; SCA-1, -2 and -6), dentatorubropallidoluysian atrophy (DRPLA) and Kennedy's disease. Specific examples of muscle degeneration include muscular dystrophy. Specific examples of cardiac degeneration include ischemia caused by heart trauma *e.g.*, associated with or resulting from a heart attack, or insufficient blood supply, or oxygen. Such disorders are but a few of the specific examples of disorders treatable with a method of the invention.

Apoptotic disorders are caused by various molecular mechanisms. For example, particular amino acid sequences are known to confer apoptosis, such as polyglutamine repeat sequences (polyGln) and protein sequences within cell receptors, such as the neurotrophin receptor. Caspases are also known to mediate the apoptotic signaling pathway in cells. Thus, conditions treatable with the compositions of the invention include cells, tissue and organs in which such apoptosis inducing or increasing proteins are expressed or active.

Bcl-B was initially identified using a genetic screen of a human liver library. A TBLASTN search of the human Expressed Sequence Tag (EST) database using the amino-acid sequence of the mouse Boo/Diva as a query identified partial cDNA having homology with Boo. A human EST clone (Accession no. AA098865) was obtained and sequenced in its entirety, revealing an open reading frame (ORF) encompassing the last 151 residues of a protein with homology to Boo (Bcl-B).

Structurally, Bcl-B is characterized as having four "BH" domains, denoted BH1, BH2, BH3 and BH4. The sequences of the domains for exemplary Bcl-B are BH1, VLSDSPGPTWGRVVTLVTFAG; BH2, AWLQAQGGWDGFCHF; BH3, EAAVLRSAARLRQI; and BH4, ERTELLADYLGYCAREPGTP (SEQ ID NOS: 3-6, respectively). These domains are present in other Bcl-2 family member proteins that inhibit apoptosis. Comparisons of this predicted sequence with all known Bcl-2 family members by

BLAST search indicated that it is most similar to the murine Bcl-2-family protein Boo (also known as Diva). Exemplary Bcl-B (SEQ ID NO:2) shares 47% amino-acid sequence identity with Boo.

The *BCL-B* gene is comprised of two exons interrupted by a ~2.3 kbp intron. The location of this intron corresponds precisely to the intronic interruption in the coding region of the anti-apoptotic *BCL-2* and *BCL-X* genes (corresponding to the motif GGW<sup>Δ</sup>D in BH2). By comparison to *BCL-B*, the pro-apoptotic genes *BAX* and *BAK* have more complicated exon-intron organizations, in which the coding regions of the gene are spread over 5 (Bak) or 6 (Bax) exons. The similar genomic organization of the *BCL-2*, *BCL-XL*, and *BCL-B* genes thus suggests they evolved from a common ancestor and indirectly implies a similar mechanism of action for their encoded proteins.

In transfection assays performed in four different human tumor cell lines, anti-apoptotic activity of Bcl-B was observed. Bcl-B binds and suppresses apoptosis-induction by Bax, but does not detectably interact with Bak or modulate apoptosis induced by Bak over-expression. However, because Bcl-B is capable of associating with either anti-apoptotic proteins (e.g., Bcl-2 and Bcl-X<sub>L</sub>) or with the pro-apoptotic proteins (e.g., Bax), it is possible that Bcl-B displays different effects on apoptosis, cell proliferation, survival, growth or differentiation depending on cellular context. A similar phenomenon has been reported for some other Bcl-2 family proteins. For example, Bcl-2 reportedly promotes apoptosis in photoreceptor cells of the retina while Bax can suppress cell death in some types of neurons (Chen et al., *Proc. Natl. Acad. Sci. USA* **93**, 7042-7047 (1996); Middleton et al., *Development* **122**, 695-701 (1996)). Thus, Bcl-B is likely to inhibit cell apoptosis, proliferation, survival, growth or differentiation, in some cellular contexts and promote cell apoptosis proliferation, survival, growth or differentiation in other cellular contexts.

The Bcl-B transmembrane (TM) domain is important for apoptosis-inhibiting activity, as well as for intracellular localization to mitochondria. Deletion of the domain reduces Bcl-B ability to inhibit apoptosis as well as reduces association with mitochondria. Replacing the native Bcl-B transmembrane domain with other transmembrane domains, including those targeting mitochondria, will likely restore Bcl-B apoptosis modulating activity. Thus, the invention includes Bcl-B polypeptides and subsequences having non-Bcl-B transmembrane domains, for example, from Bcl-2, Bax, Bak, Bcl-X<sub>L</sub>, etc.

5  
10  
15  
20  
25  
30

The correlation between membrane-targeting and function is reminiscent of some other Bcl-2 family proteins indicating that the site of action of Bcl-B is close to the intracellular organelles, including mitochondria, with which it associates. Though roughly half the Bcl-BATM protein was associated with the HM membrane fraction in cells, this may be due to its dimerization with other resident Bcl-2 family proteins. A membrane site of action for Bcl-B would be consistent with evidence that several Bcl-2 family proteins are capable of forming ion channels or pores in membranes. Indeed, molecular modeling of Bcl-B on the structure of Bcl-X<sub>L</sub> suggests that it possesses a similar overall fold and that it contains amphipathic  $\alpha$ -helices similar to the putative pore-forming  $\alpha$ 5 and  $\alpha$ 6 of Bcl-X<sub>L</sub>.

The differences observed in the activity and protein interaction partners of murine Boo and human Bcl-B proteins indicate that Bcl-B does not represent the human orthologue of mouse Boo/Diva. Also consistent with this interpretation is the difference in the expression patterns of Bcl-B and Boo. For example, although Boo (Diva) is expressed predominantly in ovary, testis, and epididymis in adult mice, Bcl-B mRNA is widely expressed in adult human tissues.

Weak interactions of Bcl-B with Apaf1 in co-immunoprecipitation assays have been detected but functional analysis has yet to reveal an effect of Bcl-B on Apaf1-induced apoptosis. Since Apaf1 is a soluble cytosolic protein, the inability of Bcl-BATM to suppress Bax-induced apoptosis suggests that Bcl-B may not play a significant role in suppressing Apaf1. Moreover, the observation that Bcl-B suppresses apoptosis induced by Bax but not Bak also argues against a role for Bcl-B as an Apaf1 suppresser, given that both Bax and Bak induce mitochondrial release of the Apaf1-activator, cytochrome c. However, though Bcl-B may not regulate Apaf1, it is possible that it binds to and suppresses the activity of some other unidentified Apaf1-like, CED4-family protein, analogous to CED9, with which Bcl-B shares 20% amino-acid sequence identity (53% similarity).

As used herein, the term “apoptosis” or “programmed cell death” means that the cell (tissue or organ) exhibits one or more characteristics associated with programmed cell death. Characteristics include inhibition of cell survival, growth, death or differentiation, protein/nucleic acid cleavage/fragmentation, chromatin condensation, membrane fragmentation, changes in expression or activity of one or more proteins that promote apoptosis or that inhibit apoptosis, for example, increased caspase (*e.g.*, caspase-3, -7 or -9) or Fas activity or expression, increased expression of proteins containing amplified polyglutamine repeat sequences or



amplification of the sequences themselves, protein aggregate formation, etc. "Modulating" apoptosis means increasing, stimulating or inducing, or decreasing, inhibiting, blocking or preventing (e.g., prophylaxis) one or more characteristics of programmed cell death as described herein or known in the art. As described herein, compositions of the invention, including Bcl-B polypeptides, antibodies and subsequences thereof and antisense can have one or more apoptosis modulating activities.

As used herein, the term "isolated," when referring to a molecule or composition, such as a nucleic acid, polypeptide or antibody of the invention, means that the molecule or composition is separated from at least one other compound, such as a protein, DNA, RNA, or other contaminants with which it is associated *in vivo* or in its naturally occurring state. Thus, a nucleic acid sequence is considered isolated when it has been isolated from any other component with which it is naturally associated. Components naturally associated with nucleic acid and protein include cellular components such as membrane, sugars, carbohydrate, lipids, fats, small molecules (e.g., hormones), etc.

As used herein, "recombinant" refers to a polynucleotide synthesized or otherwise manipulated *in vitro* (e.g., "recombinant polynucleotide"), to methods of using recombinant polynucleotides to produce gene products *in vitro*, in cells or in other biological systems, or to a polypeptide ("recombinant protein") encoded by a recombinant polynucleotide.

A composition of the invention can also be "substantially pure." For example, a substantially pure Bcl-B protein will generally be free (30% or less measured by mass) of contaminating cellular components. A substantially pure composition can be free from chemical precursors or other chemicals when chemically synthesized, such as chemical intermediates or byproducts used to synthesize proteins or nucleic acids or that are byproducts of such chemical synthesis procedures. Thus, "substantially pure" can mean Bcl-B protein having less than about 30%, 20%, 10% and more likely 5% (by dry weight), of non-Bcl-B protein (also referred to herein as a "contaminating protein"), of chemical precursors, or of culture medium, *i.e.*, culture medium is less than about 30% of the volume of the protein preparation. An isolated composition can therefore be in a homogeneous state. It can be in a dry or an aqueous solution. The invention includes isolated or purified preparations (protein or nucleic acid) of at least 0.01, 0.1, 1.0, and 10 milligrams in dry weight. Purity and homogeneity can be determined, e.g., using

analytical chemistry techniques such as, *e.g.*, polyacrylamide gel electrophoresis (SDS-PAGE) or high performance liquid chromatography (HPLC).

The terms “nucleic acid,” “nucleic acid sequence” or “polynucleotide” are used interchangeably to refer to a deoxy-ribonucleotide or ribonucleotide oligonucleotide, including single- or double-stranded forms, and coding or non-coding (*e.g.*, “antisense”) forms. Antisense includes single strand non-coding sequence complementary to coding sequence (*i.e.* sense strand), triplex forming sequences as well as single and double strand ribozymes, DNAzymes and RNAi.

The term encompasses nucleic acids containing analogues of natural nucleotides. The term also encompasses nucleic-acid-like structures with synthetic backbones. DNA backbone analogues provided by the invention include phosphodiester, phosphorothioate, phosphorodithioate, methylphosphonate, phosphoramidate, alkyl phosphotriester, sulfamate, 3'-thioacetal, methylene(methylimino), 3'-N-carbamate, morpholino carbamate, and peptide nucleic acids (PNAs); see *Oligonucleotides and Analogues, a Practical Approach*, edited by F. Eckstein, IRL Press at Oxford University Press (1991); *Antisense Strategies*, Annals of the New York Academy of Sciences, Volume 600, Eds. Baserga and Denhardt (NYAS 1992); Milligan (1993) *J. Med. Chem.* 36:1923-1937; *Antisense Research and Applications* (1993, CRC Press). PNAs contain non-ionic backbones, such as N-(2-aminoethyl) glycine units. Phosphorothioate linkages are described, *e.g.*, by U.S. Patent Nos. 6,031,092; 6,001,982; 5,684,148; see also, WO 97/03211; WO 96/39154; Mata (1997) *Toxicol. Appl. Pharmacol.* 144:189-197. Other synthetic backbones encompassed by the term include methyl-phosphonate linkages or alternating methylphosphonate and phosphodiester linkages (see, *e.g.*, U.S. Patent No. 5,962,674; Strauss-Soukup (1997) *Biochemistry* 36:8692-8698), and benzylphosphonate linkages (see, *e.g.*, U.S. Patent No. 5,532,226; Samstag (1996) *Antisense Nucleic Acid Drug Dev* 6:153-156).

The term nucleic acid is used interchangeably with gene, DNA, RNA, cDNA, mRNA, oligonucleotide primer, probe and amplification product. Nucleic acid molecules encoding Bcl-B polypeptides, proteins or amino acid sequences, hybridize to Bcl-B encoding nucleic acids or have the recited percent identity to Bcl-B set forth in SEQ ID NO:1 are collectively referred to as “nucleic acids or polynucleotides of the invention” or “Bcl-B nucleic acids or polynucleotides.”

The term “heterologous” when used in reference to a nucleic acid, indicates that the nucleic acid is in a cell or plant where it is not normally found in nature; or, comprises two or

more nucleic acid subsequences which are not found in the same relationship to each other as normally found in nature, or is recombinantly engineered so that its level of expression, or physical relationship to other nucleic acids or other molecules in a cell, or structure, is not normally found in nature. For example, a heterologous nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes, such as a recombinant chimeric protein having one or more amino acids from different proteins, arranged in a manner not found in nature; *e.g.*, a promoter sequence operably linked to a nucleic of the invention. As another example, the invention provides recombinant constructs (expression cassettes, vectors, viruses, and the like) comprising various combinations of promoters and sequences of the invention.

As used herein the terms “polypeptide,” “protein,” and “peptide” are used interchangeably and include compositions of the invention such as Bcl-B protein and antibodies that bind Bcl-B, as well as subsequences thereof. Amino acids comprise the polypeptides, all L- or D-isomers, or mixtures thereof. Amino acids also include circularized forms, such as formed by intra- or intermolecular disulfide bonds, or end-to-end amino-carboxyl linkages.

Also included are “analogs,” or “conservative variants” and “mimetics” (*e.g.*, “peptidomimetics”) with structures and activity that substantially correspond to the polypeptides of the invention, including the exemplary Bcl-B sequence as set forth in SEQ ID NO:2. Thus, the terms “conservative variant” or “analog” or “mimetic” also refer to a polypeptide or peptide which has a modified amino acid sequence, such that the change(s) do not substantially alter the polypeptide’s (the conservative variant’s) structure and/or activity (*e.g.*, to modulate apoptosis, bind to a binding molecule, etc.), as defined herein. These include conservatively modified variations of an amino acid sequence, *i.e.*, amino acid substitutions, additions or deletions of those residues that are not critical for protein activity, or substitution of amino acids with residues having similar properties (*e.g.*, acidic, basic, positively or negatively charged, polar or non-polar, etc.) such that the substitutions of even critical amino acids does not substantially alter structure and/or activity, *i.e.*, they are “substantially the same” in terms of structure or function as an unmodified sequence.

Conservative substitution tables describing chemically and structurally similar amino acids are well known in the art. For example, an exemplary guideline to select conservative substitutions includes (original residue followed by exemplary substitution): ala/gly or ser; arg/lys; asn/ gln or his; asp/glu; cys/ser; gln/asn; gly/asp; gly/ala or pro; his/asn or gln; ile/leu or val;

leu/ile or val; lys/arg or gln or glu; met/leu or tyr or ile; phe/met or leu or tyr; ser/thr; thr/ser; trp/tyr; tyr/trp or phe; val/ile or leu. An alternative uses the following six groups, each containing amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); (see also, *e.g.*, Creighton (1984) *Proteins*, W.H. Freeman and Company; Schulz and Schimer (1979) *Principles of Protein Structure*, Springer-Verlag). One of skill in the art will appreciate that the above-described substitutions are not the only possible conservative substitutions. For example, for some purposes, one may regard all charged amino acids as conservative substitutions for each other whether they are positive or negative. In addition, individual substitutions, deletions or additions that alter, add or delete a single amino acid or a small percentage of amino acids within an encoded sequence are also considered "conservatively modified variations."

A "non-essential" amino acid is a residue that can be altered from the wild-type sequence of Bcl-B (*e.g.*, the sequence of SEQ ID NO:2) without destroying or, without substantially altering a biological activity. An "essential" amino acid is defined as one which when substituted with a non-conservative amino acid or deleted results in significant loss or complete abolition of activity. For example, amino acid residues that are likely to be essential among the polypeptides of the present invention include, for example, amino acids present in one or more of the BH1, BH2, BH3 and BH4 domains. Essential amino acids are unamenable to non-conservative substitution or deletion of more than a few amino acids (*e.g.*, 1-3 amino acids).

The terms "mimetic" and "peptidomimetic" refer to a synthetic chemical compound that has substantially the same structural or functional characteristics of the polypeptides of the invention (*e.g.*, ability to modulate apoptosis, bind to Bax, Bcl-2 or Bcl-X<sub>L</sub>, associate with mitochondria, form membrane channels, etc.). The mimetic can be either entirely composed of synthetic, non-natural analogues of amino acids, or, is a chimeric molecule of partly natural peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetics' structure or activity. As with polypeptides of the invention which are conservative variants, routine tests can be used to determine whether a mimetic is within the scope of the invention, *i.e.*, that its function is not

substantially altered, *e.g.*, it retains at least part of an activity of native Bcl-B set forth in SEQ ID NO:2. Polypeptide mimetics can contain any combination of natural and non-natural structural components, which are typically from three structural groups: a) residue linkage groups other than the natural amide bond ("peptide bond") linkages; b) non-natural residues in place of naturally occurring amino acid residues; or c) residues which induce secondary structural mimicry, *i.e.*, to induce or stabilize a secondary structure, *e.g.*, a beta turn, gamma turn, beta sheet, alpha helix conformation, etc. A polypeptide can be characterized as a mimetic when one or more of its residues are joined by chemical means other than natural peptide (amide) bonds.

Individual peptidomimetic residues can be joined by peptide bonds, other chemical bonds or coupling means, such as, *e.g.*, glutaraldehyde, N-hydroxysuccinimide esters, bifunctional maleimides, N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-diisopropylcarbodiimide (DIC). Linking groups that can be an alternative to the traditional amide bond ("peptide bond") linkages include, *e.g.*, ketomethylene (*e.g.*, -C(=O)-CH<sub>2</sub>- for -C(=O)-NH-), aminomethylene (CH<sub>2</sub>-NH), ethylene, olefin (CH=CH), ether (CH<sub>2</sub>-O), thioether (CH<sub>2</sub>-S), tetrazole (CN<sub>4</sub>-), thiazole, retroamide, thioamide, or ester (see, *e.g.*, Spatola (1983) in *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, Vol. 7, pp 267-357, "Peptide Backbone Modifications," Marcell Dekker, NY).

“Expression control elements” are generally cis-acting nucleic acid transcriptional regulatory sequences. Examples include promoters capable of directing expression of a nucleic acid to cells (bacteria, plant, insect, mammalian, yeast, etc.). Examples also include enhancers capable of influencing transcription of a transcribed gene when located at a significant distance from the initiation site, or at the 5', 3' ends, or in an intron within the coding sequence. Expression control elements may contain only the minimum elements needed for transcription of the recombinant nucleic acid, such as a minimal promoter sequence.

An expression control element can be “constitutive,” such that transcription of an operably linked nucleic acid occurs without the presence of a signal or stimuli. Expression control elements can confer expression in a manner that is “regulatable,” that is, a signal or stimuli (external or internal or produced by other cells, such as a hormone produced by one cell that affects gene expression in another cell) that increases or decreases expression of the operably linked nucleic acid. A regulatable element that increases expression of the operably linked nucleic acid in response to a signal or stimuli is also referred to as an “inducible element”

operably linked nucleic acid. A regulatable element that increases expression of the operably linked nucleic acid in response to a signal or stimuli is also referred to as an “inducible element”

(i.e., is induced by a signal, e.g., a hormone). A regulatable element that decreases expression of the operably linked nucleic acid in response to a signal or stimuli is referred to as a “repressible element” (i.e., the signal decreases expression such that when the signal is removed or absent expression increases). Generally, the amount of increase or decrease conferred by such elements is proportional to the amount of signal or stimuli present. Thus, in the case of an inducible element, the greater an amount of signal or stimuli present the greater the increase in expression. Typically, basal levels of transcription are greater for a repressible element than for an inducible element.

Expression control elements and promoters include those active in a particular tissue or cell type, referred to herein as a “tissue-specific expression control elements.” Tissue-specific expression control elements are typically active in specific cell or tissue as they are recognized by transcriptional activator proteins, or other regulators of transcription unique to the specific cell or tissue type. Expression control elements also include elements that confer expression at a particular stage of the cell cycle or differentiation. Accordingly, the invention includes expression control elements that confer constitutive, regulatable, tissue-specific, cell cycle specific, and differentiation stage specific expression.

Expression control elements include other components that influence expression (transcription, translation, RNA or protein stability, etc.), and therefore, include components in addition to promoter and enhancer sequences. Such components include, for example, leader sequences and fusion partner sequences, internal ribosome binding sites (IRES) elements for the creation of multigene, or polycistronic, messages, splice signal for introns, maintenance of the correct reading frame of the gene to permit in-frame translation of mRNA, polyadenylation signal for polyadenylation of the transcript of a gene of interest, stop codons, etc. Thus, a combination of expression control elements might include a cis-acting transcriptional control element promoter, an enhancer, transcription terminator, an origin of replication, a chromosomal integration sequence, 5' and 3' untranslated regions, and an intron. Expression control elements included herein can be from bacteria, yeast, plant, insect or animal (mammalian or non-mammalian), so long as they function to control expression of an operably linked nucleic acid.

The term “expression cassette” refers to any recombinant expression system for expressing a nucleic acid sequence of the invention *in vitro* or *in vivo*, constitutively or inducibly, in any cell or organism, including, in addition to insect and plant cells, prokaryotic,

yeast, fungal or mammalian (*e.g.*, human) cells. The term includes linear or circular expression systems. The term includes all vectors. The cassettes, when present in a host cell can be episomal, that is maintained apart from the host's genome, or integrate into the genome. Expression cassettes having the ability to self-replicate or not, *i.e.*, drive only transient expression in a cell, are also included. Introduction of expression constructs into target cells can be carried out by conventional methods well known in the art (osmotic shock (*e.g.*, calcium phosphate), electroporation, viral vectors, vesicles or lipid carriers (*e.g.*, lipofection), direct microinjection, etc.

The term "operable linkage," "operably linked" and grammatical variations thereof means that the element or elements referred to are in a physical or functional relation that allows them to function in their intended manner. Thus, for an expression control element such as a promoter to be in operable linkage with a gene, the promoter modulates expression of the gene (increasing or decreasing expression, as appropriate). Likewise, enhancers operably linked to a gene influence expression of the gene, which include sequences located 5' and 3' of the initiation site and in introns, for example. Thus, nucleic acid relatively close to a gene, within a gene and flanking a gene, *e.g.*, within about 1-10, 10-50, 50-100, 100-500, 500-1000 or more nucleotides from the 5' or 3' termini are included. The invention therefore provides nucleic acids of the invention "operably linked" to an expression control element.

Another example of two elements in operable linkage is where expression of a gene influences expression of another gene. For example, a promoter driving expression of a gene encoding a transcription factor which binds to a promoter that activates expression of a second gene is operably linked to the second gene. Similarly, a Bcl-B polypeptide that binds to Bax is operably linked with a protein (or for that matter signaling pathway) influenced by Bax, through binding, or where, for example, Bax regulates transcription of a gene product. The terms operable linkage, operably linked and the like are therefore functionally defined.

The term "vector" refers to nucleic acid, such as a plasmid, virus (*e.g.*, viral vector), or other vehicle known in the art that can be manipulated by insertion or incorporation of a polynucleotide, for genetic manipulation (*i.e.*, "cloning vectors"), or can be used to transcribe or translate the inserted polynucleotide (*i.e.*, "expression vectors"). Such vectors are useful for manipulation of nucleic acids of the invention, for producing invention polypeptides (*e.g.*, baculovirus system), for introducing them into cells or whole organisms (*ex vivo* or *in vivo*), and

expressing the transcribed Bcl-B antisense or encoded Bcl-B protein in cells *in vitro* or *in vivo*.

A vector generally contains at least an origin of replication for propagation in a cell. Control elements, including expression control elements as set forth herein, present within a vector, are included to facilitate transcription.

Vectors can include a selection marker. As is known in the art, "selection marker" refers to genes that allow the selection of cells containing the gene. "Positive selection" refers to selecting the cells that survive (contain the positive selection marker) upon exposure to the positive selection agent. For example, drug resistance is a common positive selection marker; cells containing the positive selection marker will survive in culture medium containing the selection drug, and those which do not contain the resistance gene will die. Suitable drug resistance genes are *neo*, which confers resistance to G418, or *hygr*, which confers resistance to hygromycin, or *puro*, which confers resistance to puromycin. Other positive selection marker genes include genes that allow the sorting or screening of cells. These genes include genes for fluorescent proteins, the lacZ gene, the alkaline phosphatase gene, and cell surface markers such as CD8 (isolate positive cells by CD8 antibody panning), among others. "Negative selection" refers to a process whereby cells containing a negative selection marker are killed upon exposure to an appropriate negative selection agent which kills cells containing the negative selection marker. For example, cells which contain the herpes simplex virus-thymidine kinase (*HSV-tk*) gene are sensitive to the drug.

Vectors applicable in the invention are those based on viral vectors, such as simian virus 40 (SV40) or bovine papilloma virus (BPV), which has the ability to replicate as extra-chromosomal elements (Eukaryotic Viral Vectors, Cold Spring Harbor Laboratory, Gluzman ed., 1982; Sarver *et al.*, *Mol. Cell. Biol.* 1:486 (1981)). Viral vectors include retroviral (lentivirus), adeno-associated virus (see, *e.g.*, Okada (1996) *Gene Ther.* 3:957-964; Muzyczka (1994) *J. Clin. Invest.* 94:1351; U.S. Patent Nos. 6,156,303; 6,143,548 5,952,221, describing AAV vectors; see also 6,004,799; 5,833,993), adenovirus (see, *e.g.*, U.S. Patent Nos. 6,140,087; 6,136,594; 6,133,028; 6,120,764), reovirus, herpesvirus, rotavirus genomes etc., modified for introducing and directing expression of a polynucleotide or transgene in cells. Retroviral vectors can include those based upon murine leukemia virus (see, *e.g.*, U.S. Patent No. 6,132,731), gibbon ape leukemia virus (see, *e.g.*, U.S. Patent No. 6,033,905), simian immuno-deficiency virus, human immuno-deficiency virus (see, *e.g.*, U.S. Patent No. 5,985,641), and combinations thereof.



Vectors also include those that efficiently deliver genes to animal cells *in vivo* (e.g., stem cells) (see, e.g., U.S. Patent Nos. 5,821,235 and 5,786,340; Croyle, M.A. *et al.*, *Gene Ther.* 5:645 (1998); Croyle, M.A. *et al.*, *Pharm. Res.* 15:1348 (1998); Croyle, M.A. *et al.*, *Hum. Gene Ther.* 9:561 (1998); Foreman, P.K. *et al.*, *Hum. Gene Ther.* 9:1313 (1998); Wirtz, S. *et al.*, *Gut* 44:800 (1999)). Adenoviral and adeno-associated viral vectors suitable for *in vivo* delivery are described, for example, in U.S. Patent Nos. 5,700,470, 5,731,172 and 5,604,090. Additional vectors suitable for *in vivo* delivery include herpes simplex virus vectors (see, e.g., U.S. Patent No. 5,501,979), retroviral vectors (see, e.g., U.S. Patent Nos. 5,624,820, 5,693,508 and 5,674,703; and WO92/05266 and WO92/14829), bovine papilloma virus (BPV) vectors (see, e.g., U.S. Patent No. 5,719,054), CMV-based vectors (see, e.g., U.S. Patent No. 5,561,063) and parvovirus, rotavirus and Norwalk virus vectors. Lentiviral vectors are useful for infecting dividing as well as non-dividing cells (see, e.g., U.S. Patent No. 6,013,516).

Vectors for insect cell expression commonly use recombinant variations of baculoviruses and other nucleopolyhedrovirus, e.g., *Bombyx mori* nucleopolyhedrovirus vectors (see, e.g., Choi (2000) *Arch. Virol.* 145:171-177). For example, Lepidopteran and Coleopteran cells are used to replicate baculoviruses to promote expression of foreign genes carried by baculoviruses, e.g., *Spodoptera frugiperda* cells are infected with recombinant *Autographa californica* nuclear polyhedrosis viruses (AcNPV) carrying a heterologous, e.g., a human, coding sequence (see, e.g., Lee (2000) *J. Virol.* 74:11873-11880; Wu (2000) *J. Biotechnol.* 80:75-83). See, e.g., U.S. Patent No. 6,143,565, describing use of the polydnavirus of the parasitic wasp *Glyptapanteles indiensis* to stably integrate nucleic acid into the genome of Lepidopteran and Coleopteran insect cell lines. See also, U.S. Patent Nos. 6,130,074; 5,858,353; 5,004,687.

Expression vectors capable of expressing proteins in plants are well known in the art, and include, e.g., vectors from *Agrobacterium* spp., potato virus X (see, e.g., Angell (1997) *EMBO J.* 16:3675-3684), tobacco mosaic virus (see, e.g., Casper (1996) *Gene* 173:69-73), tomato bushy stunt virus (see, e.g., Hillman (1989) *Virology* 169:42-50), tobacco etch virus (see, e.g., Dolja (1997) *Virology* 234:243-252), bean golden mosaic virus (see, e.g., Morinaga (1993) *Microbiol Immunol.* 37:471-476), cauliflower mosaic virus (see, e.g., Cecchini (1997) *Mol. Plant Microbe Interact.* 10:1094-1101), maize Ac/Ds transposable element (see, e.g., Rubin (1997) *Mol. Cell Biol.* 17:6294-6302; Kunze (1996) *Curr. Top. Microbiol. Immunol.* 204:161-194), and the maize

suppressor-mutator (Spm) transposable element (see, *e.g.*, Schlappi (1996) Plant Mol. Biol. 32:717-725); and derivatives thereof.

Introduction of nucleic acid and polypeptide *in vitro*, *ex vivo* and *in vivo* can also be accomplished using other techniques. For example, a polynucleotide comprising an expression control element in operable linkage with a nucleic acid encoding Bcl-B protein can be incorporated into particles or a polymeric substance, such as polyesters, polyamine acids, hydrogel, polyvinyl pyrrolidone, ethylene-vinylacetate, methylcellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymers, polylactide/glycolide copolymers, or ethylenevinylacetate copolymers. A polynucleotide can be entrapped in microcapsules prepared by coacervation techniques or by interfacial polymerization, for example, by the use of hydroxymethylcellulose or gelatin-microcapsules, or poly (methylmethacrolate) microcapsules, respectively, or in a colloid drug delivery system. Colloidal dispersion systems include macromolecule complexes, nano-capsules, microspheres, beads, and lipid-based systems, including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The use of liposomes for introducing various compositions, including polynucleotides, is known to those skilled in the art (see, *e.g.*, U.S. Patent Nos. 4,844,904, 5,000,959, 4,863,740, and 4,975,282). A carrier comprising a natural polymer, or a derivative or a hydrolysate of a natural polymer, described in WO 94/20078 and U.S. Patent No. 6,096,291, is suitable for mucosal delivery of molecules, such as polypeptides and polynucleotides. Piperazine based amphilic cationic lipids useful for gene therapy also are known (see, *e.g.*, U.S. Patent No. 5,861,397). Cationic lipid systems also are known (see, *e.g.*, U.S. Patent No. 5,459,127). Accordingly, vector (viral and non-viral, *e.g.*, naked DNA) and non-vector means of delivery into mucosal cells or tissue, *in vitro*, *in vivo* and *ex vivo* can be achieved and are contemplated.

As used herein, the term "transgene" means a polynucleotide that has been introduced into a cell or organism by artifice. For example, in a host cell having a transgene, the transgene has been introduced by genetic manipulation or "transformation" of the cell. A cell or progeny thereof into which the transgene has been introduced is referred to as a "transformed cell" or "transformant." Typically, the transgene is included in progeny of the transformant or becomes a part of the organism (tissue or organs) that develops from the cell. Transgenes may be inserted into the chromosomal DNA or maintained as a self-replicating plasmid, YAC, minichromosome,

or the like. Transgenes include any gene that is transcribed into an antisense or encodes a polypeptide.

A “host cell” includes a cell that expresses a composition of the invention. Host cells include cultured cells (cells that have been adapted for growth *in vitro*) and primary cells isolated from an organism manipulated to contain a composition of the invention. Host cells also can be present in an animal (subject), either by *ex vivo* manipulation of cells, transgenic introduction of compositions (non-human animals), or by *in vivo* introduction of a composition of the invention.

The term “subject” refers to an animal. Typically, the animal is a mammal, however, any animal in which Bcl-B is present or may modulate apoptosis is encompassed by the term.

Specific examples are primates (humans), dogs, cats, horses, cows, pigs, and sheep. Subjects include those having a physiological disorder associated with apoptosis or at risk of developing a disorder associated with apoptosis, although they may not exhibit symptoms of the disorder. Subjects therefore may have a disorder associated with apoptosis or may not. At risk subjects can be identified by screening for genes associated with the disorders; over or under expression of a gene (*e.g.*, caspase, fas, etc.) indicates a subject at risk for a disorder.

The term “antibody” or “Ab” includes both intact antibodies having at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds and antigen binding fragments thereof, or equivalents thereof, either isolated from natural sources, recombinantly generated or partially or entirely synthetic. Examples of antigen binding fragments include, *e.g.*, Fab fragments, F(ab')<sub>2</sub> fragments, Fd fragments, dAb fragments, isolated complementarity determining regions (CDR), single chain antibodies, chimeric antibodies, partially humanized antibodies, human antibodies made in non-human animals (*e.g.*, transgenic mice) or any form of antigen binding fragment.

As used herein, the term “bind” or “binding” means that the components referred to specifically interact with each other at a molecular level. Direct binding means physical contact between the components. Indirect binding means binding to one or more components that bind. Thus, the two components need not physically contact each other in order to bind as they may be a part of an oligomeric complex in which an intermediary component binds between the two components. Indirect or direct binding may be relatively stable, such as that which occurs between an antibody and an antigen or be less stable, *e.g.* a dissociation constant ( $K_D$ ) of less than about  $10^{-6}$ . Binding may also be transient, such as the binding that occurs between a

transcription factor and DNA for transcription initiation, which does not occur in the absence of transcription. "Specific binding" is where the binding is selective between the components.

Specific binding can be detected using methods known in the art, for example, by immunoprecipitation, affinity chromatography, gel shift assays, gene expression assays, etc.

"Specific" and "non-specific" binding can be distinguished using appropriate controls.

The terms "array," "microarray," "DNA array" or "nucleic acid array" or "biochip" as used herein mean a plurality of target elements, each target element comprising a defined amount of one or more biological molecules (*e.g.*, nucleic acid, protein, drugs, small organic compounds, natural compounds, libraries thereof, etc.), including the Bcl-B nucleic acids or proteins of the invention, immobilized on a substrate (*e.g.*, solid or semi-solid, permeable or non-permeable surface) for binding with sample nucleic acids, proteins, antibodies, test agents, drugs, organic or natural compounds. The Bcl-B proteins, nucleic acids, antibodies of the invention and subsequences thereof can be incorporated into any form of microarray, such as those described, *e.g.*, in U.S. Patent Nos. 6,045,996; 6,022,963; 6,013,440; 5,959,098; 5,856,174; 5,770,456; 5,556,752; 5,143,854.

The term "plant" includes whole plants, plant parts (*e.g.*, leaves, stems, flowers, roots, etc.), plant protoplasts, seeds and plant cells and progeny of same, including progeny obtained through cuttings or through regeneration of a plant or plant tissue from a plant cell. The class of plants which can be used in the method of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including angiosperms (monocotyledonous and dicotyledonous plants), as well as gymnosperms. It includes plants of a variety of ploidy levels, including polyploid, diploid, haploid and hemizygous states. Plantlets are also included within the meaning of "plant." Suitable plants for use in the invention include any plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants. Examples of monocotyledonous plants include, but are not limited to, asparagus, field and sweet corn, barley, wheat, rice, sorghum, onion, pearl millet, rye and oat, and ornamentals. Examples of dicotyledonous plants include, but are not limited to, tomato, potato, arabidopsis, tobacco, cotton, rapeseed, field beans, soybeans, peppers, lettuce, peas, alfalfa, clover, cole crops or Brassica (*e.g.*, cabbage, broccoli, cauliflower, brussel sprouts), radish, carrot, beets, eggplant, spinach, cucumber, squash, melons, cantaloupe, sunflowers and various ornamentals. The term "plant cell", as used herein, refers to protoplasts, gamete producing cells, and cells that are

capable of regenerating into whole plants. Accordingly, a seed comprising multiple plant cells capable of regenerating into a whole plant is included in the definition of "plant cell". As used herein, "plant tissue" includes differentiated and undifferentiated tissues of a plant, including but not limited to roots, stems, shoots, leaves, pollen, seeds, tumor tissue and various forms of cells and culture such as single cells, protoplast, embryos, and callus tissue.

A "biotic insult," as used herein, refers to plant challenge caused by viable or biologic agents (biotic agents), such as insects, fungi, bacteria, viruses, nematodes, viroids, mycoplasmas, and the like.

An "abiotic insult," as used herein, refers to plant challenge by a non-viable or non-living agent (abiotic agent). Abiotic agents that can cause challenge include, for example, environmental factors such as low moisture (drought), high moisture (flooding), nutrient deficiency, radiation levels, air pollution (ozone, acid rain, sulfur dioxide, etc.), temperature (hot and cold extremes), and soil toxicity, as well as herbicide damage, pesticide damage, or other agricultural practices (e.g., over-fertilization, improper use of chemical sprays, etc.).

The term "pathogen" refers to any biological organism or chemical agent produced by a biological organism that causes a disease or disease state in an animal or plant, including, but not limited to viruses, fungi, bacterium, nematodes, and other related microorganisms.

"Functional subsequences," which are portions of the compositions of the invention that retain at least one activity or function, all or in part, of the native full-length composition are included. "Functional subsequences" therefore include portions of Bcl-B protein (e.g., fragments of SEQ ID NO:2) which modulate apoptosis, bind to Bax, Bcl-2 and Bcl-X<sub>L</sub>, associate with mitochondria or form a membrane channel alone or in combination with other Bcl-2 protein family members. Functional subsequences also include subsequences that are immunogenic.

Bcl-B protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of SEQ ID NO:2, which include less amino acids than the full length Bcl-B, and exhibit all or part of at least one activity of a Bcl-B protein (e.g., apoptosis modulating activity). Typically, an active portion comprises a domain or motif with at least one activity of the Bcl-B protein, for example, a domain or motif capable of modulating apoptosis; a domain or motif capable of interacting with a binding molecule, such as itself, Bax, Bcl-2 or Bcl-X<sub>L</sub>; a domain or motif capable of associating with mitochondria, such as Bcl-B transmembrane domain; a domain or motif capable of modulating an intracellular molecule (e.g.,

Bax) that participates in an apoptotic signaling pathway; a domain or motif capable of forming a membrane channel alone, or in combination with other Bcl-2 protein family members; a domain or motif capable of modulating a caspase; a domain or motif capable of modulating DNA fragmentation, chromatin condensation or other processes associated with apoptosis; or a domain or motif capable of modulating cell proliferation, survival, growth or differentiation.

A biologically active portion of a Bcl-B protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length. Biologically active portions can be used directly as agents that modulate apoptosis or other activities of full length Bcl-B. Bcl-B protein can also be used as targets for developing agents which modulate a Bcl-B mediated activity, *e.g.*, an activity described herein, and are potential therapeutics for treating disorders associated with undesirable or insufficient/deficient apoptosis.

The terms percent "sequence identity" or "sequence homology," in the context of two or more nucleic acids or polypeptide sequences, refers to sequences or subsequences thereof that have a specified percentage of nucleotides (or amino acid residues) that are the same, when compared and aligned for maximum correspondence over a comparison window. Percent identity can be measured by manual alignment and visual inspection or using a sequence comparison algorithm, as described herein. This definition also refers to the complement (antisense strand) of a sequence.

For example, in various embodiments, polypeptides of the invention include those having an amino acid sequence at least about 70%, at least about 80%, at least about 90%, at least about 95% and at least about 99% identical to an exemplary sequence set forth in SEQ ID NO:2. Polypeptide subsequences are also included. In additional embodiments, a subsequence is at least about 15, 20, 25, 30, 40, 50, 75, 125, 150 or 200, or greater amino acids (*e.g.*, 300, 350, 400, 450, 500, 550, etc.) in length. Thus, a polypeptide sequence having the requisite sequence identity to SEQ ID NO:2, or a subsequence thereof, also is a polypeptide of the invention. In various additional embodiments, the polypeptides, including subsequences have one or more activities of a sequence set forth in SEQ ID NO:2. Thus, in various aspects a polypeptide has apoptosis modulating activity, oligomerization (heteromerizes or forms homodimers, trimers, etc.) activity, binds to Bcl-2, B

Nucleic acids of the invention include those having at least about 70%, at least about 80%, at least about 90%, at least about 95%, and at least about 99% identity to the exemplary

sequence set forth in SEQ ID NO:1, wherein the sequence is distinct from Accession no. AA098865. Thus, if a nucleic acid sequence has the requisite sequence identity to SEQ ID NO:1, or a subsequence thereof, it also is a polynucleotide sequence of the invention. In various aspects, the sequence is less than about 50Kb, 25 Kb, 10Kb, 5Kb or 2.5Kb. In additional aspects, the sequence is between about 2.5-1.0Kb, 1.0-0.5Kb, 0.25-0.1 Kb and 100-15 base pairs. In yet other aspects, the sequence is selected from SEQ ID NO:1; SEQ ID NO:1 where one or more T's are U's; nucleic acids complementary to these sequences and subsequences of the aforementioned sequences at least 15 base pairs long.

In one aspect, the percent identity exists over a region of the sequence that is at least about 25 nucleotides or amino acid residues in length, or, over a region that is at least about 50 to 100 nucleotides or amino acids in length. Parameters (including, e.g., window sizes, gap penalties and the like) to be used in calculating "percent sequence identities" between two nucleic acids or polypeptides to identify and determine whether one is within the scope of the invention are described in detail, below.

As used herein, the term "hybridizes under stringent conditions" describes conditions for hybridization and washing. Stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Hybridization stringency typically depends on the stringency of the wash. A particular example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 50°C. Another example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 55°C. A further example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C. Highly stringent conditions include one or more washes in 0.2X SSC, 0.1% SDS at 65°C following the hybridization.

Sequences included in the invention are those that hybridize to exemplary Bcl-B sequence set forth in Example 1 (SEQ ID NO:1). Such sequences may be the entire length of the sequence, or may be fragments. For example, in various embodiments, the sequence that hybridizes has a length of about 12-30, 30-50, 50-100, 100-250, 250-500, 500-1000, 1000-2500,

2500-5000 or 5000-10000 base pairs. Such sequences are also distinct from Accession no. AA098865.

Such sequences are useful as probes, for example, to detect the presence or amount of Bcl-B in a sample; to detect a disorder associated with apoptosis where under or over expression of Bcl-B is indicative of the disorder (or indicative of a predisposition to the disorder); for identifying other Bcl-B family genes in mammalian (*e.g.*, human) or other organisms, *e.g.*, bacteria, plant, insect, yeast, etc. Such sequences are also useful as antisense. For example, sequences that hybridize to double strand Bcl-B (*e.g.*, genomic Bcl-B) form a triplex nucleic acid molecule thereby inhibiting expression of the target Bcl-B. Double stranded RNA sequences (known as "RNAi") for inhibiting gene expression in organisms are also applicable and are known in the art (Kennerdell *et al.*, *Cell* 95:1017-1026 (1998); Fire *et al.*, *Nature*, 391:806-811 (1998)). Such antisense sequences can interfere with Bcl-B protein activity or expression and in turn increase apoptosis, or decrease cell survival or proliferation.

The invention further provides chimeric polypeptides comprising an amino acid sequence of the invention, for example, a sequence having at least about 65% identity to SEQ ID NO:2, and a second polypeptide sequence. Thus, peptides and polypeptides of the invention are synthesized and expressed as chimeric or "fusion" proteins with one or more additional domains linked thereto for, *e.g.*, to more readily isolate or identify a recombinantly synthesized peptide, and the like. Detection and purification facilitating domains include, *e.g.*, metal chelating peptides such as polyhistidine tracts and histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle WA). The inclusion of a cleavable linker sequences such as Factor Xa or enterokinase (Invitrogen, San Diego CA) between the purification domain and GCA-associated peptide or polypeptide can be useful to facilitate purification. For example, an expression vector can include an epitope-encoding nucleic acid sequence linked to six histidine residues followed by a thioredoxin and an enterokinase cleavage site (see, *e.g.*, Williams (1995) *Biochemistry* 34:1787-1797; Dobeli (1998) *Protein Expr. Purif.* 12:404-14). The histidine residues facilitate detection and purification while the enterokinase cleavage site provides a means for purifying the epitope from the remainder of the fusion protein.



Polypeptides and peptides of the invention can be isolated from natural sources, be synthetic, or be recombinantly generated polypeptides. Peptides and proteins can be recombinantly expressed *in vitro* or *in vivo*. The peptides and polypeptides of the invention can be made and isolated using any method known in the art.

Polypeptide and peptides of the invention can also be synthesized, whole or in part, using chemical methods well known in the art. See *e.g.*, Caruthers (1980) Nucleic Acids Res. Symp. Ser. 215-223; Horn (1980) Nucleic Acids Res. Symp. Ser. 225-232; Banga, A.K., Therapeutic Peptides and Proteins, Formulation, Processing and Delivery Systems (1995) Technomic Publishing Co., Lancaster, PA. For example, peptide synthesis can be performed using various solid-phase techniques (see *e.g.*, Roberge (1995) Science 269:202; Merrifield (1997) Methods Enzymol. 289:3-13) and automated synthesis may be achieved, *e.g.*, using the ABI 431A Peptide Synthesizer (Perkin Elmer). The skilled artisan will recognize that individual synthetic residues and polypeptides incorporating mimetics can be synthesized using a variety of procedures and methodologies, which are well described in the scientific and patent literature, *e.g.*, Organic Syntheses Collective Volumes, Gilman, et al. (Eds) John Wiley & Sons, Inc., NY. Polypeptides incorporating mimetics can also be made using solid phase synthetic procedures, as described, *e.g.*, by Di Marchi, et al., U.S. Pat. No. 5,422,426. Peptides and peptide mimetics of the invention can also be synthesized using combinatorial methodologies. Various techniques for generation of peptide and peptidomimetic libraries are well known, and include, *e.g.*, multipin, tea bag, and split-couple-mix techniques; see, *e.g.*, al-Obeidi (1998) Mol. Biotechnol. 9:205-223; Hruby (1997) Curr. Opin. Chem. Biol. 1:114-119; Ostergaard (1997) Mol. Divers. 3:17-27; Ostresh (1996) Methods Enzymol. 267:220-234. Modified peptides of the invention can be further produced by chemical modification methods, see, *e.g.*, Belousov (1997) Nucleic Acids Res. 25:3440-3444; Frenkel (1995) Free Radic. Biol. Med. 19:373-380; Blommers (1994) Biochemistry 33:7886-7896.

The invention further provides isolated or recombinant nucleic acids comprising a nucleic acid sequence having at least about 80% sequence identity to SEQ ID NO:1, further including expression cassettes (*e.g.*, vectors), cells and transgenic animals comprising the nucleic acids of the invention. As the genes and vectors of the invention can be made and expressed *in vitro* or *in vivo*, the invention provides for a variety of means of making and expressing these genes and vectors. One of skill will recognize that desired phenotypes associated with altered gene activity

can be obtained by modulating the expression or activity of the genes and nucleic acids (*e.g.*, promoters) within the expression cassettes (*e.g.*, vectors) of the invention. Any of the known methods described for increasing or decreasing expression or activity can be used for this invention. The invention can be practiced in conjunction with any method or protocol known in the art, which are well described in the scientific and patent literature.

The nucleic acid sequences of the invention and other nucleic acids used to practice this invention, whether RNA, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed recombinantly. Any recombinant expression system can be used, including, in addition to insect and bacterial cells, *e.g.*, mammalian, yeast or plant cell expression systems.

Alternatively, these nucleic acids can be synthesized *in vitro* by well-known chemical synthesis techniques, as described in, *e.g.*, Belousov (1997) Nucleic Acids Res. 25:3440-3444; Frenkel (1995) Free Radic. Biol. Med. 19:373-380; Blommers (1994) Biochemistry 33:7886-7896; Narang (1979) Meth. Enzymol. 68:90; Brown (1979) Meth. Enzymol. 68:109; Beaucage (1981) Tetra. Lett. 22:1859; U.S. Patent No. 4,458,066.

Techniques for the manipulation of nucleic acids, such as, *e.g.*, generating mutations in sequences, subcloning, labeling probes, sequencing, hybridization and the like are well described in the scientific and patent literature, see, *e.g.*, Sambrook, ed., MOLECULAR CLONING: A LABORATORY MANUAL (2ND ED.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989); CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); LABORATORY TECHNIQUES IN BIOCHEMISTRY AND MOLECULAR BIOLOGY: HYBRIDIZATION WITH NUCLEIC ACID PROBES, Part I. Theory and Nucleic Acid Preparation, Tijssen, ed. Elsevier, N.Y. (1993).

Expression control elements, which, are operably linked to the coding sequence for the exemplary polypeptide of the invention, SEQ ID NO:2 (or antisense), including promoters, comprising the genomic sequences 5' (upstream) of a transcriptional start site (see SEQ ID NO:1), enhancers and intronic sequences are readily available in the art or can be obtained using routine molecular biological techniques (*e.g.*, cell expression assays). Methods for further characterization of promoter sequences include those general methods described by, *e.g.*, Pang (1997) Biotechniques 22:1046-1048; Gobinda (1993) PCR Meth. Applic. 2:318; Triglia (1988) Nucleic Acids Res. 16:8186; Lagerstrom (1991) PCR Methods Applic. 1:111; Parker (1991)

Nucleic Acids Res. 19:3055. As is apparent to one of ordinary skill in the art, these techniques can also be applied to identify, characterize and isolate any expression control sequences corresponding to or associated with the nucleic acid and polypeptide sequences of the invention.

The invention provides oligonucleotide primers that can amplify all or any specific region within a nucleic acid sequence of the invention, particularly, the exemplary SEQ ID NO:1. The nucleic acids of the invention can also be mutated, detected, generated or measured quantitatively using amplification techniques. Using the nucleic acid sequences of the invention (e.g., as in the exemplary SEQ ID NO:1), the skilled artisan can select and design suitable oligonucleotide amplification primers. Amplification methods are also known in the art, and include, e.g., polymerase chain reaction, PCR (see, e.g., PCR PROTOCOLS, A GUIDE TO METHODS AND APPLICATIONS, ed. Innis, Academic Press, N.Y. (1990) and PCR STRATEGIES (1995), ed. Innis, Academic Press, Inc., N.Y.); ligase chain reaction (LCR) (see, e.g., Barringer (1990) Gene 89:117); transcription amplification (see, e.g., Kwoh (1989) Proc. Natl. Acad. Sci. USA, 86:1173); and, self-sustained sequence replication (see, e.g., Guatelli (1990) Proc. Natl. Acad. Sci. USA, 87:1874); Q Beta replicase amplification (see, e.g., Smith (1997) J. Clin. Microbiol. 35:1477-1491; Burg (1996) Mol. Cell. Probes 10:257-271) and other RNA polymerase mediated techniques (e.g., NASBA, Cangene, Mississauga, Ontario).

The invention provides transformed cells comprising a nucleic acid of the invention. The cells can be mammalian (such as human), insect (such as *Spodoptera frugiperda*, *Spodoptera exigua*, *Spodoptera littoralis*, *Spodoptera litura*, *Pseudaletia separata*, *Trichopusia ni*, *Plutella xylostella*, *Bombyx mori*, *Lymantria dispar*, *Heliothis virescens*, *Autographica californica* and other insect, particularly lepidopteran and coleopteran, cell lines), plant, bacteria, yeast, etc. Techniques for transforming and culturing cells are well described in the scientific and patent literature; see, e.g., Weiss (1995) Methods Mol. Biol. 39:79-95, describing insect cell culture in serum-free media; Tom (1995) Methods Mol. Biol. 39:203-224; Kulakosky (1998) Glycobiology 8:741-745; Altmann (1999) Glycoconj. J. 16:109-123; Yanase (1998) Acta Virol. 42:293-298; U.S. Patent Nos. 6,153,409; 6,143,565; 6,103,526.

The invention also provides transgenic animals, including non-human mammals and insects. Insects stably expressing the nucleic acids of the invention can be to study apoptosis, screening for modulators of apoptosis and caspases, manipulation of insect life cycles, such as *Bombyx mori* and its use in silk production. The nucleic acids of the invention can be expressed

in a variety of insect larvae, *e.g.*, *Bombyx mori* (see, *e.g.*, Maeda (1985) Nature 315: 592-594), *Trichoplusia ni*, the cabbage looper larvae (Medin (1990) Proc. Nat. Acad. Sci. USA 87: 2760-2764) and *Manduca sexta*, the tobacco hornworm (U.S. Patent No. 5,471,858). See, *e.g.*, Keshan (2000) J. Insect Physiol. 46:1061-1068; U.S. Patent No. 5,118,616.

5 Non-human transgenic mammals include, *e.g.*, primates (chimps, apes, macaques, orangutans), cows, goats, horses, sheep, pigs, rats, rabbits and mice, comprising the nucleic acids of the invention. These animals can be used, *e.g.*, as *in vivo* models to study apoptosis and potential therapies for apoptosis. For example, animal models that mimic human apoptotic disorders (*e.g.*, cell degeneration or proliferation, such as that of neural or muscular tissue, or cancer), can be used to study whether Bcl-B compositions of the invention ameliorate (relieve or improve one or more symptoms, *e.g.*, inhibit additional cell death or degeneration) of the disorder. Such models can be used to screen for agents that modulate Bcl-B activity *in vivo*. Transgenic non-human animals can be designed and generated using any method known in the art; see, *e.g.*, U.S. Patent Nos. 6,156,952; 6,118,044; 6,111,166; 6,107,541; 5,959,171; 5,922,854; 5,892,070; 5,880,327; 5,891,698; 5,639,940; 5,573,933, describing making and using transgenic mice, rats, rabbits, sheep, pigs and cows. See also, *e.g.*, Pollock (1999) J. Immunol. Methods 231:147-157, describing the production of recombinant proteins in the milk of transgenic dairy animals; Baguisi (1999) Nat. Biotechnol. 17:456-461, demonstrating the production of transgenic goats.

20 The invention also provides transgenic plants, including seeds, expressing the Bcl-B nucleic acids and polypeptides of the invention. Nucleic acids may be introduced into the genome of the desired plant host by a variety of conventional techniques. For example, the DNA construct may be introduced directly into the genomic DNA of the plant cell using techniques such as electroporation and microinjection of plant cell protoplasts, or the DNA constructs can be introduced directly to plant tissue using ballistic methods, such as DNA particle bombardment. Microprojectile bombardment to deliver DNA into plant cells is an alternative means of transformation for the numerous species considered recalcitrant to *Agrobacterium*- or protoplast-mediated transformation methods. For example, see, *e.g.*, Christou (1997) Plant Mol. Biol. 35:197-203; Pawlowski (1996) Mol. Biotechnol. 6:17-30; Klein (1987) Nature 327:70-73; 25 Takumi (1997) Genes Genet. Syst. 72:63-69, discussing use of particle bombardment to introduce transgenes into wheat; and Adam (1997) *supra*, for use of particle bombardment to 30

introduce YACs into plant cells. For example, Rinehart (1997) *supra*, used particle bombardment to generate transgenic cotton plants. Apparatus for accelerating particles is described U.S. Pat. No. 5,015,580; and, the commercially available BioRad (Biolistics) PDS-2000 particle acceleration instrument; see also, U.S. Patent Nos. 5,608,148 and 5,681,730, describing particle-mediated transformation of gymnosperms.

Alternatively, transformed plant cells can be generated by fusion of the recipient cells with bacterial protoplasts containing DNA, use of DEAE dextran, polyethylene glycol precipitation, as described, *e.g.*, in Paszkowski (1984) EMBO J. 3:2717-2722. DNA construct can be introduced directly into the genomic DNA of the plant cell using electroporation, as described, *e.g.*, in Fromm (1985) Proc. Natl. Acad. Sci. USA 82:5824, or by microinjection of plant cell protoplasts, as described, *e.g.*, Schnorf (1991) Transgenic Res. 1:23-30.

DNA can also be introduced in to plant cells using recombinant viruses. Plant cells can be transformed using viral vectors, such as, *e.g.*, tobacco mosaic virus derived vectors (Rouwendal (1997) Plant Mol. Biol. 33:989-999), see Porta (1996) "Use of viral replicons for the expression of genes in plants," Mol. Biotechnol. 5:209-221.

Alternatively, the DNA constructs may be combined with suitable T-DNA flanking regions and introduced into a conventional *Agrobacterium tumefaciens* host vector. The virulence functions of the *Agrobacterium tumefaciens* host will direct the insertion of the construct and adjacent marker into the plant cell DNA when the cell is infected by the bacteria.

*Agrobacterium tumefaciens*-mediated transformation techniques, including disarming and use of binary vectors, are well described in the scientific literature. See, *e.g.*, Horsch (1984) *Science* 233:496-498; Fraley (1983) *Proc. Natl. Acad. Sci. USA* 80:4803 (1983); *Gene Transfer to Plants*, Potrykus, ed. (Springer-Verlag, Berlin 1995).

The nucleic acid sequences of the invention include genes and gene products identified and characterized by analysis using the exemplary protein and nucleic acid sequences of the invention, including SEQ ID NO:2 and SEQ ID NO:1. For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters are used unless alternative parameters are designated herein. The sequence comparison algorithm then calculates the percent sequence

identity for the test sequence(s) relative to the reference sequence, based on the designated or default program parameters. A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 25 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Methods of sequence alignment for comparison are well-known in the art. Optimal alignment of sequences can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (CLUSTAL, GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection.

In one aspect of the invention, a CLUSTAL algorithm, such as the CLUSTAL W program, is used to determine if a nucleic acid or polypeptide sequence is within the scope of the invention; see, *e.g.*, Thompson (1994) *Nuc. Acids Res.* 22:4673-4680; Higgins (1996) *Methods Enzymol* 266:383-402. Variations of this aspect can also be used, such as CLUSTAL X, see Jeanmougin (1998) *Trends Biochem Sci* 23:403-405; Thompson (1997) *Nucleic Acids Res* 25:4876-4882. CLUSTAL W program, described by Thompson (1994) *supra*, in the methods of the invention used with the following parameters: K tuple (word) size: 1, window size: 5, scoring method: percentage, number of top diagonals: 5, gap penalty: 3.

Another algorithm is PILEUP, which can be used to determine whether a polypeptide or nucleic acid has sufficient sequence identity to SEQ ID NO:1 or SEQ ID NO:2 to be with the scope of the invention. This program creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The following parameters are used

with PILEUP in the methods of the invention: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps.

Another example of an algorithm suitable for determining percent sequence identity (*i.e.*, substantial similarity or identity) in the invention is the BLAST algorithm, which is described in Altschul (1990) *J. Mol. Biol.* 215:403-410. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul (1990) *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues, always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. In one embodiment, to determine if a nucleic acid sequence is within the scope of the invention, the BLASTN program (for nucleotide sequences) is used incorporating as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as default parameters a wordlength (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix (see, *e.g.*, Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915).

The invention provides antibodies that specifically bind to the Bcl-B polypeptides of the invention, *e.g.*, the exemplary SEQ ID NO:2. These antibodies can be used, *e.g.*, to isolate the polypeptides of the invention, to identify the presence of Bcl-B polypeptides that modulate apoptosis, and the like. To generate antibodies, Bcl-B polypeptides or peptides (antigenic fragments of SEQ ID NO:2) can be conjugated to another molecule or can be administered with an adjuvant. The coding sequence can be part of an expression cassette or vector capable of expressing the immunogen *in vivo* (see, *e.g.*, Katsumi (1994) *Hum. Gene Ther.* 5:1335-9).

1007174-00000

Methods of producing polyclonal and monoclonal antibodies are known to those of skill in the art and described in the scientific and patent literature, see, e.g., Coligan, CURRENT PROTOCOLS IN IMMUNOLGY, Wiley/Greene, NY (1991); Stites (eds.) BASIC AND CLINICAL IMMUNOLOGY (7th ed.) Lange Medical Publications, Los Altos, CA; Goding, MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE (2d ed.) Academic Press, New York, NY (1986); Harlow (1988) ANTIBODIES, A LABORTAROY MANUAL, Cold Spring Harbor Publications, New York.

Antibodies also can be generated *in vitro*, e.g., using recombinant antibody binding site expressing phage display libraries, in addition to the traditional *in vivo* methods using animals. See, e.g., Huse Science 246:1275 (1989); Ward Nature 341:544 (1989); Hoogenboom Trends Biotechnol. 15:62-70 (1997); Katz Annu. Rev. Biophys. Biomol. Struct. 26:27-45 (1997). Human antibodies can be generated in mice engineered to produce only human antibodies, as described by, e.g., U.S. Patent Nos. 5,877,397; 5,874,299; 5,789,650; and 5,939,598. B-cells from these mice can be immortalized using standard techniques (e.g., by fusing with an immortalizing cell line such as a myeloma or by manipulating such B-cells by other techniques to perpetuate a cell line) to produce a monoclonal human antibody-producing cell. See, e.g., U.S. Patent Nos. 5,916,771 and 5,985,615.

As discussed, the invention includes Bcl-B polypeptides having apoptosis modulating activity, including subsequences and sequence variants (e.g., a polypeptide having at least about 65% identity to SEQ ID NO:2), including mimetics etc., having apoptosis modulating activity.

In addition, the invention provides methods for identifying an agent that modulates an activity of a polypeptide of the invention (e.g., a Bcl-B polypeptide having apoptosis modulating activity) that is capable inhibiting apoptosis in a cell.

Determining the amount or degree of apoptosis in a cell can be measured using routine *in vitro* or cell based techniques described herein or known in the art. For example, the amount or degree of activity can be determined by expressing a test Bcl-B polypeptide in a cell and measuring apoptosis in the cell (e.g., see Example 1 where percent apoptotic cells is determined by counting cells having nuclear fragmentation or chromatin condensation). Caspase protease activity in a cell generally reflects apoptosis and, therefore, increasing caspase activity or expression indicates increased cell apoptosis. Additional assays include measuring the amount or degree of DNA fragmentation in a cell; the amount or degree of cleavage of caspase substrates in a cell; or by measuring the amount or degree of any surrogate marker of apoptosis in the cell.



*In vitro* and cell based assay can be performed alone or in association with other pro- or anti-apoptotic proteins. Methods for measuring apoptosis are described, for example, in Methods in Enzymology (2000) volume 322, edited by John C. Reed, Academic Press, *e.g.*, chapters on pages 3, 15, 41, describing assays to measure apoptosis and surrogate markers of apoptosis and enzymes with activity related to levels of apoptosis, *e.g.*, assays to determine DNA fragmentation, caspase assays, measuring annexin V (see, *e.g.*, Zhang (1997) Biotechniques 23:525-531), and the like. See, *e.g.*, van Engeland (1996) Cytometry 24:131-139; Gorczyca (1998) Methods Mol. Biol. 91:217-238. See also, *e.g.*, U.S. Patent Nos. 6,165,737; 6,165,732; 6,160,095; 6,143,522; 6,087,384; 6,077,684; 6,060,238; 6,054,436; 5,985,829; 5,976,786; 5,952,189. The invention incorporates all such methods and variations thereof.

The invention further provides kits comprising one or more compositions of the invention, including pharmaceutical compositions. In one embodiment, a kit contains one or more Bcl-B polypeptides or polynucleotide sequences encoding Bcl-B, and a label or packaging insert including instructions for expressing the polypeptide in cells (*in vitro*, *ex vivo* or *in vivo*) under conditions allowing increased expression of a target gene, in suitable packaging material. In another embodiment, a kit contains a Bcl-B antisense or ribozyme or DNAzyme or RNAi, and a label or packaging insert including instructions for expressing or introducing the antisense or ribozyme or DNAzyme or RNAi in cells (*in vitro*, *ex vivo* or *in vivo*) under conditions allowing decreased expression Bcl-B in the cell, in suitable packaging material.

As used herein, the term "packaging material" refers to a physical structure housing the components of the kit, such as Bcl-B polypeptides, polynucleotides and antibodies. The packaging material can maintain the components sterily, and can be made of material commonly used for such purposes (*e.g.*, paper, corrugated fiber, glass, plastic, foil, etc.). The label or packaging insert can include appropriate instructions, for example, practicing a method of the invention, *e.g.*, treating a apoptotic disorder treatable by expressing a Bcl-B protein, antisense or ribozyme or DNAzyme or RNAi or antibody *in vivo*.

The invention also provides methods of treating disorders associated with apoptosis or with disorders in which apoptotic proteins participate. Disorders associated with apoptosis include disorders characterized by abnormal or undesirable, or increased or undesirable cell death, degeneration or insufficient/deficient or increased or undesirable cell growth, proliferation, survival or differentiation. Disorders in which apoptotic proteins participate are

non-apoptotic disorders, but expression or activity of proteins associated with the disorder may exacerbate the disorder.

Examples of cellular proliferative or differentiative disorders include cancer, *e.g.*, carcinoma, sarcoma, metastatic disorders or hematopoietic neoplastic disorders, *e.g.*, leukemias.

5 A metastatic tumor can arise from a multitude of primary tumor types, including but not limited to those of breast, colon, lung, and liver origin.

As used herein, the terms “cancer”, “hyperproliferative” and “neoplastic” refer to cells having the capacity for autonomous growth, *i.e.*, an abnormal state or condition characterized by rapidly proliferating cell growth. Hyperproliferative and neoplastic disease states may be categorized as pathologic, *i.e.*, characterizing or constituting a disease state, or may be categorized as non-pathologic, *i.e.*, a deviation from normal but not associated with a disease state. The term is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. “Pathologic hyperproliferative” cells occur in disease states characterized by malignant tumor growth. Examples of non-pathologic hyperproliferative cells include proliferation of cells associated with wound repair.

The terms “cancer” or “neoplasms” include malignancies of the various organ systems, such as affecting lung, breast, thyroid, lymphoid, gastrointestinal, and genito-urinary tract, as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer and/or testicular tumors, non-small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus.

25 The term “carcinoma” is art recognized and refers to malignancies of epithelial or endocrine tissues including respiratory system carcinomas, gastrointestinal system carcinomas, genitourinary system carcinomas, testicular carcinomas, breast carcinomas, prostatic carcinomas, endocrine system carcinomas, and melanomas. Exemplary carcinomas include those forming from tissue of the cervix, lung, prostate, breast, head and neck, colon, liver and ovary. The term also includes carcinosarcomas, *e.g.*, which include malignant tumors composed of carcinomatous and sarcomatous tissues. An “adenocarcinoma” refers to a carcinoma derived from glandular tissue or in which the tumor cells form recognizable glandular structures.

30 Additional examples of proliferative disorders include hematopoietic neoplastic disorders. As used herein, the term “hematopoietic neoplastic disorders” includes diseases

5  
10  
15  
20  
25  
30

involving hyperplastic/neoplastic cells of hematopoietic origin, *e.g.*, arising from myeloid, lymphoid or erythroid lineages, or precursor cells thereof. Typically, the diseases arise from poorly differentiated acute leukemias, *e.g.*, erythroblastic leukemia and acute megakaryoblastic leukemia. Additional exemplary myeloid disorders include, but are not limited to, acute promyeloid leukemia (APML), acute myelogenous leukemia (AML) and chronic myelogenous leukemia (CML) (reviewed in Vaickus, L. (1991) Crit Rev. in Oncol./Hematol. 11:267-97); lymphoid malignancies include, but are not limited to acute lymphoblastic leukemia (ALL) which includes B-lineage ALL and T-lineage ALL, chronic lymphocytic leukemia (CLL), prolymphocytic leukemia (PLL), hairy cell leukemia (HLL) and Waldenstrom's macroglobulinemia (WM). Additional forms of malignant lymphomas include, but are not limited to non-Hodgkin lymphoma and variants thereof, peripheral T cell lymphomas, adult T cell leukemia/lymphoma (ATL), cutaneous T-cell lymphoma (CTCL), large granular lymphocytic leukemia (LGF), Hodgkin's disease and Reed-Sternberg disease.

Degenerative disorders include neuro-or muscular degeneration or atrophy. Such disorders afflict the brain and include, for example, Alzheimer's disease, Parkinson's disease, Creutzfeldt-Jacob's disease (CJD), Huntington disease (HD), Machado-Joseph disease (MJD), Spinocerebellar ataxias 1, 2 and 6 (SCA-1, -2 and -6), dentatorubropallidoluysian atrophy (DRPLA), Kennedy's disease, ischemia, stroke and head trauma.

Treatment generally results in reducing or preventing the severity or symptoms of the disorder in the subject, *i.e.*, an improvement in the subject's condition or a "therapeutic effect." Therefore, treatment can reduce the severity or prevent one or more symptoms of the apoptotic disorder, inhibit progression or worsening of the disorder, and in some instances, reverse the disorder. Thus, in the case of a degenerative disorder, for example, treatment can prevent further degeneration resulting in a stabilization of the condition. Improvement in a cell proliferative disorder can mean inhibition of tumor growth, at least in part, such as to slow the progression of the tumor. This can also lead to an improvement in the subject because slowing the growth of a tumor generally results in prolonging life span of the patient or reducing the chances for metastasis, even though the subject may not be free of the tumor.

As used herein, the term "ameliorate" means an improvement in the subject's condition, a reduction in the severity of a disorder, or an inhibition of progression or worsening of the disorder.

10 The doses or "effective amount" for treating a subject are preferably sufficient to  
ameliorate one, several or all of the symptoms of the condition, to a measurable or detectable  
extent, although preventing or inhibiting a progression or worsening of the disorder or condition,  
or a symptom, is a satisfactory outcome. Thus, to ameliorate a disorder treatable by a method of  
5 the invention, that is, a disorder where modulating apoptosis results in an improvement, the  
effective amount will depend on the condition and the desired outcome and can be readily  
ascertained by the skilled artisan. Appropriate amounts will also depend upon the therapeutic  
effect desired, as well as the individual subject (e.g., the bioavailability within the subject,  
gender, age, etc.). The effective amount can be ascertained by measuring relevant physiological  
effects.

15 The methods of the invention for treating a subject are applicable for prophylaxis to  
prevent an apoptotic condition in a subject, such as a degenerative condition or an associated  
disorder. The methods of the invention for treating a subject also can be supplemented with  
other forms of therapy. Supplementary therapies include drug treatment (e.g., growth factor  
addition, chemotherapy, radiation treatment, surgical resection, etc.). The skilled artisan can  
readily ascertain therapies that may be used in combination with the treatment methods of the  
invention.

20 As the methods of the invention include methods of treating a subject, such as a human  
subject, the invention also provides "pharmaceutically acceptable" or "physiologically  
acceptable" formulations including one or more compositions of the invention alone, in  
combination with each other, or in combination with other drugs or agents suitable for treatment.  
Such formulations can be administered *ex vivo* or *in vivo* to a subject in order to practice the  
treatment methods of the invention, for example.

25 As used herein, the terms "pharmaceutically acceptable" and "physiologically  
acceptable" refer to carriers, diluents, excipients and the like that can be administered to a  
subject, preferably without producing excessive adverse side-effects (e.g., nausea, abdominal  
pain, headaches, etc.). Such preparations for administration include sterile aqueous or non-  
aqueous solutions, suspensions, and emulsions.

30 Pharmaceutical formulations can be made from carriers, diluents, excipients, solvents,  
dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying  
agents, and the like, compatible with administration to a subject. Such formulations can be

contained in a tablet (coated or uncoated), capsule (hard or soft), microbead, emulsion, powder, granule, crystal, suspension, syrup or elixir. Supplementary active compounds and preservatives, among other additives, may also be present, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

5 A pharmaceutical formulation can be formulated to be compatible with its intended route of administration. Thus, pharmaceutical formulations include carriers, diluents, or excipients suitable for administration by routes including intraperitoneal, intradermal, subcutaneous, oral (e.g., ingestion or inhalation), intravenous, intracavity, intracranial, transdermal (topical), parenteral, e.g. transmucosal and rectal.

Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

20 Pharmaceutical formulations suitable for injection include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid  
25 polyethylene glycol, and the like), and suitable mixtures thereof. Fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. Isotonic agents, for  
30 example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride can be included in the

composition. Prolonged absorption of injectable formulations can be achieved by including an agent that delays absorption, for example, aluminum monostearate or gelatin.

For oral administration, a composition can be incorporated with excipients and used in the form of tablets, troches, or capsules, *e.g.*, gelatin capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included in oral formulations. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or flavoring.

Formulations can also include carriers to protect the composition against rapid degradation or elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. For example, a time delay material such as glyceryl monostearate or glyceryl stearate alone, or in combination with a wax, may be employed.

Additional formulations include biodegradable or biocompatible particles or a polymeric substance such as polyesters, polyamine acids, hydrogel, polyvinyl pyrrolidone, polyanhydrides, polyglycolic acid, ethylene-vinylacetate, methylcellulose, carboxymethylcellulose, protamine sulfate, or lactide/glycolide copolymers, polylactide/glycolide copolymers, or ethylenevinylacetate copolymers in order to control delivery of an administered composition. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc., for example.

The rate of release of a composition can be controlled by altering the concentration or composition of such macromolecules. For example, the composition can be entrapped in microcapsules prepared by coacervation techniques or by interfacial polymerization, for example, by the use of hydroxymethylcellulose or gelatin-microcapsules or poly (methylmethacrylate) microcapsules, respectively, or in a colloid drug delivery system. Colloidal dispersion systems include macromolecule complexes, nano-capsules, microspheres, microbeads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles,

and liposomes. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

Additional pharmaceutical formulations appropriate for administration are known in the art and are applicable in the methods and compositions of the invention (see, e.g., Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing Co., Easton, PA; The Merck Index (1996) 12th ed., Merck Publishing Group, Whitehouse, NJ; and Pharmaceutical Principles of Solid Dosage Forms, Technomic Publishing Co., Inc., Lancaster, Pa., (1993)).

Injection needles, endoscopes, cannulas, intubation tubes, catheters and the like can be used to deliver the formulation to various parts of a subject. This allows effective delivery and targeting of vectors to particular areas within a subject.

All publications, patents and other references cited herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. Unless defined otherwise herein, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention pertains.

As used herein, the singular forms "a", "and," and "the" include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to "a transformed cell" includes a plurality of such cells and reference to "an isolated or recombinant polynucleotide" includes reference to one or more such sequences, and so forth.

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, the following examples are intended to illustrate but not limit the scope of invention described in the claims.

### Example 1

This example describes the initial cloning of exemplary Bcl-B set forth in SEQ ID NO:1.

This example also describes routine techniques for characterizing and analyzing Bcl-B activity.

#### Cloning of BCL-B cDNAs

A TBLASTN search of the human Expressed Sequence Tag (EST) database using the amino-acid sequence of the mouse Boo/Diva as a query resulted in the identification of homologous partial cDNAs. A human EST clone (Accession no. AA098865) was obtained (Research Genetics) and sequenced in its entirety, revealing an open reading frame (ORF)

1007741520702

encompassing the last 151 residues of a protein with homology to Boo (Bcl-B). The corresponding genomic sequence for this cDNA was identified in the human genome database (clone CTD-2184D3), which was derived from human chromosome 15q21.

Because the EST clone lacked a candidate start codon, the corresponding 5'-end of Bcl-B cDNAs was cloned by a reverse-transcriptase polymerase chain reaction (RT-PCR) approach, using the forward primer NK01 18 (5'-CGGGCCAAGAAAACCAGCGAAGG-3'; SEQ ID NO:7), which was designed to hybridize to a region upstream of the Bcl-B ORF as predicted from the genomic data, and the reverse primer NK0121 (5'-CACTCAAGGAAGAGCCATTTCAT-3'; SEQ ID NO:8), which is complementary to a region downstream of the predicted Bcl-B ORF corresponding to the 3'-untranslated region of the putative mRNA. PCR amplification using template human liver cDNA (Clontech) with the above primers yielded a single ~ 0.9 Kb product which was cloned into pCR2.1-TOPO (Invitrogen, following manufacturer's instructions) to generate TOPO-Bcl-B (pNK254) and sequenced. The cDNA containing the complete open reading frame (ORF) corresponding to a 204 amino-acid protein is shown in Figure 1A.

Plasmid construction

The ORF encoding Bcl-B was PCR-amplified from TOPO-Bcl-B (pNK254) using the forward primer NK0101 (5'-GGAATTCATGGTTGACCAGTTGCGGGAG-3'; SEQ ID NO:9) and reverse primer NK0103 (5'-CCGCTCGAGTCATAATAATCGTGTCCAGAG-3'; SEQ ID NO:10). The PCR products were digested with *Eco*R1 and *Xho*I and cloned into the *Eco*R1 and *Xho*I sites of pcDNA3-Myc (Stratagene), and the *Eco*R1 and *Sa*I sites of pcl-Neo-Flag (Invitrogen) and pEGFP-C2 (Clontech). A plasmid encoding Bcl-B lacking its C-terminal transmembrane domain (Bcl-BΔTM) was constructed by PCR-based mutagenesis using primers NK0101 and NK0131 (5'-CCGCTCGAGTCATGTTTCTCCAAAAGCCAGTG-3'; SEQ ID NO:11). The resulting PCR product was digested with *Eco*R1 and *Xho*I and cloned into pcDNA3-Myc.

RT-PCR analysis

Expression of *BCL-B* mRNA in various tissues was examined by RT-PCR, using oligo-dT primed first-strand cDNA derived from multiple adult human tissues (Clontech) as templates. cDNAs were amplified following manufacturer's instructions using the forward primer NK0120



(5'-GTGGTGACGCTCGTGACCTTCG-3'; SEQ ID NO:12) and NK0121 as the reverse primer. G3PDH primers were used as a positive control (Kitada et al., Antisense Res. Dev. 4, 71-79 (1994)).

#### Cell culture, transfection, and apoptosis assays

HEK293, Cos7, HT1080 and PPC1 cells were maintained in Dulbecco's modified Eagle's medium (Irvine Scientific) supplemented with 10% fetal bovine serum, 1 mM L-glutamine, and antibiotics. For apoptosis assays, cells (5X105) in 6-well dishes were co-transfected using Superfect (Qiagen) with 0.5 µg pcDNA3-Bax plus 0.5 µg of green fluorescence protein (GFP) marker plasmid pEGFP (CLONTECH) or 0.5 µg pEGFP-Bak, and 1 µg of pcDNA3, pcDNA3-Myc-Bcl-B, pcDNA3-Myc-Bcl-BΔTM, or pcDNA3-Flag-Bcl-XL. The total amount of DNA was 3 µg per transfection using pcDNA3. At 24 hrs post-transfection, both adherent and floating cells were collected, fixed and stained with 0.1 µg/ml 4',6-diamidino-2-phenylindole (DAPI). The percent of apoptotic cells were determined by counting the GFP-positive cells having nuclear fragmentation and/or chromatin condensation (Mean ± S.D.; n = 3).

#### Co-immunoprecipitation and immunoblotting assays

293T cells (5X10<sup>6</sup>) cultured with 50 µM benzoyl-Val-Ala-Asp-fluoromethylketone (ZVAD-fmk, Bachem) were co-transfected with 1.5 µg of pcDNA3-Myc-Bcl-B, pcDNA3-Flag-Bcl-B, pcDNA3-hCBP (Human Calyculin Binding Protein) (used as a control), or pcDNA3-Flag-BCI-X<sub>L</sub>, together with 1.5 µg of pEGFP, pEGFP-Bcl-B, pcDNA3-HA-BAG1, pcDNA3-HA-Bax, pcDNA3-Flag-Bcl-X<sub>L</sub>, pRC-CMV-Bcl-2 or pEGFP-Bak. At 24 hrs post-transfection, cells were collected and resuspended in lysis buffer (142.4 mM KCl, 5 mM MgCl<sub>2</sub>, 10 mM HEPES (pH7.4), 0.5 mM EGTA, 0.2% NP-40) containing 12.5 mM p-glycerophosphate, 2 mM NaF, 1 mM Na<sub>3</sub>VO<sub>4</sub>, 1 mM DTT, 1 mM PMSF and a protease inhibitor mixture (Boehringer Mannheim). Soluble lysates were incubated with 10 µl anti-Myc (Santa Cruz) or anti-Flag (Sigma) antibody conjugated Sepharose beads overnight at 4°C. Beads were washed four times in 1.5 ml lysis buffer, and boiled in Laemmli gel-loading solution before performing SDS-PAGE/immunoblotting. The following polyclonal or monoclonal antibodies were used: polyclonal rabbit anti-GFP (Roche), monoclonal rat anti-HA (Roche), monoclonal mouse anti-Flag (Sigma), monoclonal mouse anti-Myc (Santa Cruz), rabbit anti-huBcl-2, or rabbit anti-

huBcl-XL (Krajewski, et al., Am. J. Pathol. 145, 515-525 (1994); Krajewski et al., Cancer Res. 54, 5501-5507 (1994)).

### Immunofluorescence and subcellular fractionation

For immunofluorescence analysis, Cos7 cells were transfected with pEGFP-Bcl-B. At 24 hrs post-transfection, 50 uM Mitotracker Red CMXRos (Molecular Probes) was added to the cell cultures for 30 mins. The cells were washed with PBS and fixed with 4% formaldehyde before imaging by confocal microscopy using a Bio-Rad MRC 1024 instrument.

For subcellular fractionation experiments, 293T cells ( $2 \times 10^6$ ) were transfected with 10 ug pcDNA3-Myc-Bcl-B, pcDNA3-Myc-Bcl-BATM, or pRC-CMV-Bcl-2. At 24 hrs post-transfection, cells were collected and lysed by homogenization in hypotonic buffer (5 mM Tris-Cl, pH7.4, 5 mM KCl, 1.5 mM MgCl<sub>2</sub>) containing 1 mM PMSF and a mixture of other protease inhibitors (Boehringer Mannheim). The nuclear fraction was discarded by centrifugation at 500xg for 5 mins. The heavy membrane (HM) fraction was obtained by centrifugation of the post-nuclear lysates at 10,000xg for 20 mins. The resulting supernatant was either used directly or further centrifuged at 150,000xg for 90 mins to separate the light membrane fraction (LM) (pellet) from the cytosolic (C) (supernatant) fraction. Fractions were normalized to cell equivalents and analyzed by immunoblotting using antibodies specific for Myc-epitope tag (Santa Cruz) and Bcl-2, with Enhanced chemiluminescence (ECL) detection (Amersham Pharmacia Biotech).

### **Example 2**

This example describes structural characteristics of Bcl-B sequences. This example also describes expression patterns of Bcl-B in adult human tissue.

Exemplary Bcl-B nucleic acid sequence (SEQ ID NO:1) is as follows:

```
cgggccaaga aaaccagcga aggcccgccg cccacgcaga ggcgggacca
tggttgacca gttgcgggag cgcaccacca tggccgaccc gctgctgggag
cgcacccgagc tgttgctggc cgactacctg ggttactgcg cccgggaacc
cggcaccccc gagccggcgc catccacgcc cgaggccgcc gtgctgcgct
ccgcggccgc caggttacgg cagattcacc ggtccttttt ctccgcctac
ctcggttacc ccgggaaccg cttcgagctg gtggcgctga tggcggattc
cgtgctctcc gacagccccc gcccccacctg gggcagagtg gtgacgctcg
```

tgaccttcgc agggacgctg ctggagagag ggccgctggt gaccgcccgg  
 tgaagaagt ggggcttcca gccgcggcta aaggagcagg agggcgacgt  
 cggccgggac tgccagcgcc tgggtggcctt gctgagctcg cggctcatgg  
 ggcagcacgg cgcttgctg caggctcagg gcggctggga tggcttttgt  
 caactcttca ggacccctt tccactggct ttttgagaa aacagctggt  
 ccaggctttt ctgtcatgct tgtaacaac agccttcatt tatctctgga  
 cagcattatt atgagtttta aaacttttaa cccgcttcta cctgcccaac  
 tgtgaccaac taaatgacag atgtgtgaga acaagaactg agggaaagca  
 ccttcccca cccagacgt ttttatctga atgcataca ggagtcctga  
 ggtggtgatt tggccagtgt tttaacttgt gacaagtact cagggtgtgag  
 gacaagaatg caaatggctc ttccttgagt gaaagaa

The predicted ORF was initiated by a AUG start codon within a favorable Kozak context. The protein contains regions resembling the BH1, BH2, BH3, and BH4 domains typical of anti-apoptotic members of the Bcl-2 family, as well as a putative carboxyl-terminal TM domain (Figure 1B). Comparisons of this predicted sequence with all known Bcl-2 family members by BLAST search indicated that it is most similar to the murine Bcl-2-family protein Boo (also known as Diva; Song et al., *EMBO J* 18, 167-178 (1999); Inohara et al., *J. Biol. Chem.* 273, 32479-32486(1998)). Exemplary Bcl-B shares 47% amino-acid sequence identity with Boo.

The BCL-B gene is located on chromosome 15 (map 15q21), as determined by in silico screening of the human genome database at NCBI. Comparison of the BCL-B cDNA sequence with genomic data indicates a two exon structure, in which the region encoding residues Trp163 and Asp164 (within the BH2 domain) of the protein are interrupted by an ~2.3 kb intron.

Adult human tissues were analyzed for Bcl-B expression. In brief, first-strand cDNAs were made from RNA samples from adult human tissues and PCR-amplified using BCL-B-specific primers. The reverse primer was downstream of the intron to avoid amplification of contaminating genomic DNA. PCR products were size-fractionated by electrophoresis in 1% agarose gels, stained with ethidium bromide, and photographed under UV-illumination. Primers specific for glyceraldehyde-3-phosphate dehydrogenase (G3PDH) were also used for PCR as a positive control. The results indicate that BCL-B mRNA is widely expressed in adult human tissues (Figure 1C). Greatest levels of expression were detected in liver, pancreas and spleen.

### Example 3

This example describes studies demonstrating that Bcl-B binds with itself as well as other Bcl-2 family proteins. This example also describes studies demonstrating that Bcl-B binding is selective for other proteins associated with the apoptotic signaling pathway.

In brief, 293T cells were transiently transfected with various combinations of plasmids encoding Myc-Bcl-B, Myc-hCBP, GFP, GFP-Bcl-B, GFP-Bak, Bcl-2, Flag-Bcl-X<sub>L</sub>, HA-Bax, HA-BAG1 and Flag-Bcl-B. Cell lysates were prepared and immunoprecipitated as described above using either anti-Myc or anti-Flag monoclonal antibodies.

Immunoprecipitated proteins were fractionated by SDS-PAGE and subjected to immunoblot analysis using rabbit polyclonal antibodies specific for GFP, Bcl-X<sub>L</sub>, or Bcl-2 or rat monoclonal antibody specific for HA (Figure 2, top panel). To confirm expression of all proteins, equivalent volumes of lysates were also loaded directly in gels and analyzed by SDS-PAGE/immunoblotting (Figure 2, middle and bottom panels) using antibodies specific for GFP, HA, Flag, Myc, Bcl-X<sub>L</sub>, or Bcl-2. For efficiency of presentation, only the portion of the gels containing the relevant bands is shown. Additional controls, including immunoprecipitations using negative-control antibodies were also performed.

These results indicate that Bcl-B is capable of binding with itself, Bax, Bcl-2, and Bcl-X<sub>L</sub>, but not with Bak. These findings are consistent with Bcl-B suppressing Bax mediated cellular apoptosis, described in detail below.

### Example 4

This example describes data showing that Bcl-B inhibits apoptosis induced by Bax. This example also describes data showing that Bcl-B binding to Bax, at least in part, contributes to inhibiting apoptosis induced by Bax, and that Bcl-B does not detectably inhibit apoptosis induced by Bak.

The anti-apoptotic activity of Bcl-B was studied by transient transfection of a variety of cell lines, including HEK293T, Cos7 (monkey kidney cells), HT1080 and PPC1. In brief, HEK293T (Figure 3A) and Cos-7 (Figure 3B) cells at ~50% confluence in 6-well dishes were co-transfected with plasmids encoding GFP (0.5 ug) (used as a marker for transfection with Bax) and 0.5 ug of pcDNA3 (control), pcDNA3-Bax or pEGFP-Bak in combination with 1 ug (two-fold excess) of pcDNA3, pcDNA3-Myc-Bcl-B or pcDNA3-Flag-Bcl-X<sub>L</sub>. After 24 hrs

posttransfection, cells were collected and stained with DAPI. The percentage of GFP-positive cells with apoptotic morphology (fragmented nuclei or condensed chromatin) was determined (mean + SD; n = 3). Immunoblotting of control cultures supplemented with 50  $\mu$ M zVAD-fmk to prevent apoptosis confirmed production of all proteins (as described in Example 3). Similar results were also obtained with HT1080 and PPC1 cells.

Bcl-B over-expression did not induce apoptosis in any of the cell lines. Bcl-B also did not suppress apoptosis caused by over-expressing Bcl-2 or Bcl-X<sub>L</sub>. These data therefore indicate that Bcl-B is not a proapoptotic protein (induces or stimulates apoptosis) in these particular cell lines.

The cytoprotective activity of Bcl-B was studied by ascertaining the effect of Bcl-B expression on apoptosis induced by the pro-apoptotic proteins Bax and Bak. Co-expressing Bcl-B markedly suppressed apoptosis induced by Bax but not Bak (Figure 3), thus correlating the protein-binding data indicating that Bcl-B binds Bax but not Bak (Figure 2). Bcl-B mediated apoptotic suppression was not due to reduced levels of Bax protein, as determined by immunoblotting. In contrast to Bcl-B, co-expression of Bcl-X<sub>L</sub> suppressed apoptosis induced by either Bax or Bak (Figure 3). Thus, Bcl-B appears to be functionally distinct from Bcl-X<sub>L</sub>.

Binding studies confirmed that binding between Bcl-B and Bax mediates suppression of apoptosis by Bax. In brief, HEK293T cells were transiently transfected with several Myc-tagged constructs (human calcyclin-binding protein, wildtype Bcl-B, L86A, R96Q, FFR(169-171)AAA,  $\Delta$ 118-133) and cell lysates were prepared and immunoprecipitated with anti-Myc antibody conjugated to beads. Immunoprecipitates were subjected to SDS-PAGE/immunoblot analysis using rabbit polyclonal antibodies specific for Bax (Figure 4A). To confirm expression of proteins, lysates were subject to immunoblot analysis using antibodies specific for Bax and Myc.

For caspase assays, 293T cells were co-transfected with 0.5  $\mu$ g of Bax and 1.0  $\mu$ g of various Bcl-B constructs. Cell lysates were prepared 24 hrs after transfection, normalized for protein contents (25  $\mu$ g), and incubated with 100  $\mu$ M DEVD-AFC. Enzyme activity was determined by the release of AFC-fluorescence (Figure 4B).

The results in Figure 4 indicate that Bcl-B binding to Bax (Bcl-B wild type and FFR mutant) correlate with decreased caspase activity, whereas the three Bcl-B mutants L86A, R96Q and  $\Delta$ 118-133 that failed to bind Bax did not alter caspase activity (compare Myc-Bcl-B to Bcl-B

and myc-FFR to FFR169 in Figures 4A and 4B). This data confirms that binding of Bcl-B to Bax mediates the protective activity of Bcl-B in Bax mediated apoptosis.

### Example 5

This example describes data showing that Bcl-B associates with mitochondria. This example also describes data showing that the C-terminal transmembrane (TM) domain of Bcl-B is important for efficient membrane targeting and function.

A plasmid encoding Green Fluorescent Protein (GFP)-tagged Bcl-B (GFP-Bcl-B) was transfected into Cos-7 cells as described in Example 1. Two-color confocal microscopy revealed a punctate cytosolic pattern and partial co-localization of GFP-Bcl-B with a mitochondria-specific dye (MitoTracker) (Figure 5A). In contrast, cells transfected with GFP control protein produced diffuse cellular fluorescence.

To further evaluate location of Bcl-B, 293T cells were transfected with pcDNA3-Myc-Bcl-B, pcDNA3-Myc-Bcl-BATM, or pRC-CMV-Bcl-2 as described in Example 1. Subcellular fractionation analysis revealed that the Myc-tagged Bcl-B protein resides predominantly in the mitochondria-containing heavy-membrane (HM) fraction, similar to Bcl-2, as determined by immunoblot analysis of the cellular fractions (Figures 5B,C). In contrast to full-length Bcl-B, a truncation mutant of Bcl-B lacking the carboxylterminal TM domain (Bcl-BATM) targeted less efficiently to the HM fraction (Figure 5D). The Bcl-BATM protein also was ineffective at suppressing Bax-induced apoptosis (Figure 5E), even though this protein was produced at comparable levels to the full-length Bcl-B protein. Thus, efficient organellar targeting appears to be required for the apoptosis inhibiting activity of Bcl-B.

For Figure 5, at 24 hr post-transfection, cells were collected, and post-nuclear lysates prepared ("total"). An aliquot of these lysates was then fractionated by differential centrifugation at 10,000xg to pellet heavy membranes (HM). The resulting supernatant ("sup") was then either analyzed directly or subjected to centrifugation at 150,000xg to pellet light membranes (LM) and achieve a cytosolic supernatant. Proteins from each fraction were normalized relative to cell numbers and subjected to SDS-PAGE/immunoblot analysis using antibodies specific for Myc or Bcl-2. Figure 5E, 293T cells were co-transfected with a plasmid encoding GFP (used as a marker for transfection) and either pcDNA3 (control; CNTL) or

pcDNA3-Bax, in combination with a 2-fold excess of pcDNA3 (control), pcDNA3Myc-Bcl-B or pcDNA3-Myc-Bcl-BΔTM. Cells were collected and stained with DAPI after 24 hrs.

Expression of Green Fluorescent Protein (GFP)-tagged Bcl-B in cells revealed a punctate cytosolic pattern and partial co-localization with a mitochondria-specific dye (MitoTracker), as determined by two-color confocal microscopy (Figure 5A). Crude subcellular fractionation analysis revealed that Myc-tagged Bcl-B protein resides predominantly in the mitochondria-containing heavy-membrane (HM) fraction, similar to Bcl-2, as determined by immunoblot analysis of the cellular fractions (Figures 5B,C). In contrast to full-length Bcl-B, a truncation mutant of Bcl-B lacking the carboxylterminal TM domain (Bcl-BΔTM) targeted less efficiently to the HM fraction (Figure 5D). The Bcl-BΔTM protein also was ineffective at blocking Bax-induced apoptosis (Figure 5E), even though this protein was produced at comparable levels to the full-length Bcl-B protein. Thus, efficient organellar targeting appears to be required for optimal function of Bcl-B.

These results demonstrate that Bcl-B associates with mitochondria. This is not an unexpected finding given that many Bcl-2 family proteins associate with mitochondria in cells (Reed, J. *Oncogene* **17**, 3225-3236(1998); Reed, J. C. *Amer J Pathol* **157**, 1415-1430(2000)).

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.